

PERSPECTIVES IN BASIC SCIENCE

Polymorphism of host response genes: Implications in the pathogenesis and treatment of acute renal failure

**BERTRAND L. JABER, BRIAN J.G. PEREIRA, JOSEPH V. BONVENTRE,
and VAIDYANATHAPURAM S. BALAKRISHNAN**

Division of Nephrology, Caritas St. Elizabeth's Medical Center, Boston, Massachusetts; Division of Nephrology, Tufts-New England Medical Center, Boston, Massachusetts; and Renal Division, Brigham and Women's Hospital, Boston, Massachusetts

Polymorphism of host response genes: Implications in the pathogenesis and treatment of acute renal failure. Acute inflammatory disorders are the result of an interaction between genetic and environmental factors, and are often characterized by an imbalance between pro- and anti-inflammatory host immune responses. Over the past decade, polymorphisms of host response genes have been explored as genetic risk and prognostic markers in the course and severity of acute inflammatory disorders.

Increasing evidence supports an important role for inflammatory mechanisms in the pathogenesis of acute renal failure (ARF) following both ischemic and nephrotoxic injury. The use of genetic epidemiology may become a useful tool to identify patients with an altered susceptibility to developing hospital-acquired ARF, and stratify those who may benefit from preventive therapies that modulate host immune responses in a favorable way.

This review summarizes the existing experimental and clinical studies supporting the role of inflammation in ARF and critically appraises studies that have examined polymorphism of immune response genes as potential determinants of susceptibility to and severity of acute inflammatory disorders. Conclusions are drawn on the application of genetic epidemiology to the field of ARF and the rationale for further research on the role of genetic markers in ARF.

Acute renal failure (ARF) is a serious complication that develops in hospitalized patients, with an incidence of 5% to 10% [1], and case fatality rates often exceed 50% [2]. These alarming figures have not changed in decades despite tremendous advances in supportive and dialysis care, and they have been ascribed in part to increasing age and a high prevalence of comorbid conditions. ARF has also been independently associated with increased risk of

death following radiocontrast procedures [3] and following cardiac surgery [4]. Interventions to prevent or improve treatment of ARF are therefore urgently needed.

There is mounting evidence to support the critical role of inflammation in the pathogenesis of ARF. This review article is based on the premise that acute inflammatory disorders are the result of an interaction between genetic and environmental factors. Over the past decade, genetic studies of common disorders have shed some light on the relative importance of genetic risk and prognostic markers in acute and chronic inflammatory disorders.

This review summarizes the experimental and clinical data supporting the role of inflammation in ARF and critically appraises studies that have examined the importance of genetic markers as potential determinants of susceptibility to and severity of acute inflammatory disorders. Conclusions are drawn on the application of genetic epidemiology to the field of ARF.

HUMAN GENETIC VARIABILITY

Although a person's genotype represents the blending of parental genotypes, two unrelated persons share over 99.9% of their DNA sequences [5]. Many efforts both in the academic and commercial sector are under way to categorize and catalog the variations observed in the remaining 0.1% of the human genome [6]. These variants, known as gene polymorphisms, are markers of biologic diversity, and some genotypic variations correlate with specific phenotypes relevant to human disease [7, 8]. These emerging research efforts are based on the premise that human gene polymorphisms may be instrumental in developing the future practice of "genomic medicine" and individualized health care [6]. However, it is not clear whether many of these variants are incriminated in the pathogenesis of diseases, or are merely in proximity to other pathogenic genetic factors, a phenomenon known as linkage disequilibrium.

Polymorphism of human genes may be observed at one or more of the following sites (Fig. 1): (1) the promoter

Key words: acute renal failure, critical illness, infection, sepsis, inflammation, cytokines, chemokines, toll-like receptors, heat shock proteins, mortality.

Received for publication February 27, 2004
and in revised form May 13, 2004
Accepted for publication July 9, 2004

© 2005 by the International Society of Nephrology




Fig. 1. Structure of a human gene, sites of polymorphism, and functional relevance. (Modified from [205]; reprinted with permission. Jaber BL et al: *Blood Purif* 22: 101–111, 2004).

or 5'-flanking region; (2) the exon(s) or the gene coding sequences; (3) the intron(s) or the gene intervening sequences; and (4) the 3'-untranslated (3'-UTR) region.

Polymorphism involving the promoter region (5'-flanking region) of the gene may affect transcriptional activity and can therefore be of functional relevance [9]. Polymorphism of the coding region or exon may be silent or affect gene expression or function by resulting in changes in the structure, binding or trafficking of the protein. Introns are transcribed into RNA but are cut out of the messenger RNA (mRNA) before it is translated into a protein molecule. Consequently, polymorphism involving the intron may lead to defects in RNA and mRNA processing. Finally, polymorphism in the 3'-UTR region may affect gene expression by affecting RNA half-life or influencing ribosomal translation of mRNA [10].

TYPES OF HUMAN GENE POLYMORPHISM

Three types of human gene polymorphisms have been described [9] (Fig. 2): (1) single nucleotide polymorphism (SNP); (2) variable number of tandem repeats (VNTR) or minisatellite polymorphism; and (3) microsatellite polymorphism.

SNP is the most common class and normally consists of single nucleotide substitution (Fig. 2A). SNP in the promoter region of a gene may affect transcription factor binding, transcriptional control, or other aspects of gene expression.

Minisatellite polymorphism or VNTR results from the insertion in tandem of multiple copies of a nucleotide sequence of less than 100 bp long, known as a minisatellite, between two restriction sites (Fig. 2B). This class of polymorphism is characterized by many alleles, based on the number of minisatellite copies [9].

Microsatellites are stretches of DNA in which a short motif of one to five nucleotides is repeated several times

(Fig. 2C). A dinucleotide repeat of cytosine and adenine (CA)_n is the most commonly described form. Microsatellites show heritable and stable differences among individuals and are therefore polymorphic.

IDENTIFICATION OF GENES RELEVANT TO HUMAN DISEASE

There are two approaches to identifying genes relevant to human disease: linkage analyses and association studies. Linkage analyses identify coinheritance of a specific phenotype with a region of the genome in families. These analyses are successful in identifying novel genes for "monogenic" diseases, and environmental factors tend to play a minimal role in disease expression. One such example is polycystic kidney disease. Association studies identify susceptibility genes for common "polygenic" diseases. This "candidate gene" approach relies on biologic plausibility to help identify highly likely candidate genes. In this analytical approach, gene environment interactions play a critical role in disease expression. Examples would include the study of genes encoding for cytokines in sepsis and ARF.

We will next review host immune responses in acute inflammatory disorders and specifically, the inflammatory network in ARF, with special emphasis on selected pathophysiologic pathways incriminated in sepsis-, ischemic-, and nephrotoxic-mediated injury.

THE HOST IMMUNE RESPONSE IN ACUTE INFLAMMATORY STATES

During an acute inflammatory state, the host immune response is often governed by a balance of two opposing forces, a proinflammatory response geared toward recruitment of inflammatory cells to sites of injury and an anti-inflammatory response aimed at limiting tissue injury and promoting healing. These immune responses




Fig. 2. Types of gene polymorphism. (A) Single nucleotide polymorphism. (B) Variable number of tandem repeats or minisatellite loci polymorphism (DNA sequence 10 to 100 bp in length). (C) Microsatellite loci polymorphism (DNA sequence of repeating units of 2 to 4 nucleotides).

are normally regulated by monocytes and lymphocytes. In bacterial sepsis, a very common acute inflammatory state, this balance has been termed the systemic inflammatory response syndrome (SIRS) and the compensatory anti-inflammatory response syndrome (CARS), and these cellular responses are usually driven by monocytes and neutrophils. The following is a summary of these emerging concepts.

The systemic inflammatory response syndrome (SIRS)

The SIRS is an exaggerated acute host defense response to different triggering factors such as gram-negative bacterial endotoxin, trauma, and burns [11]. This leads to the release of biologically active mediators, which activate several inflammatory cascades leading to organ dysfunction, including ARF [12]. In the United States,

sepsis, an example of SIRS, is increasingly identified as a common cause of morbidity and mortality, especially in the intensive care unit setting [13].

Cytokines act in concert with specific cytokine inhibitors and soluble cytokine receptors to regulate the host response. During an acute inflammatory state, stimuli cause monocytes to release such cytokines as tumor necrosis factor- α (TNF- α), interleukin-1 β (IL-1 β) and interleukin-6 (IL-6), which in turn cause widespread vascular endothelial cell injury. Indeed, TNF- α and IL-1 β interact with the vascular endothelium, and stimulate platelet-activating factor, prostanooids, and nitric oxide synthesis, resulting in arterial vasodilatation, hypotension, and organ dysfunction. Furthermore, interleukin-8 (IL-8), a potent chemokine, facilitates the recruitment of neutrophils to inflammatory sites, which in turn, release their toxic cellular contents, including reactive oxygen

species (ROS) and proteolytic enzymes. IL-6 also induces the hepatic synthesis of positive acute-phase proteins, such as C-reactive protein (CRP) and suppresses synthesis of negative acute-phase proteins, such as albumin [14]. TNF- α and IL-6 also contribute to a series of metabolic events leading to protein catabolism, a state of negative nitrogen balance and significant loss of lean body mass [15]. In recent years, substantial evidence has accumulated linking proinflammatory cytokines to adverse outcomes among patients with acute inflammatory states [16, 17].

The compensatory anti-inflammatory response syndrome (CARS)

The CARS is believed to play a critical role in counterbalancing the proinflammatory responses associated with the SIRS. This secondary immune response is characterized by the production of immunomodulatory molecules that control the proinflammatory cytokine response, thereby, counterbalancing the SIRS. Their critical role in systemic inflammatory states is becoming recognized [18]. Major anti-inflammatory monocyte-derived molecules include IL-10, IL-1 receptor antagonist (IL-1Ra), and soluble TNF receptors (sTNF-R).

IL-10 is the most potent anti-inflammatory cytokine, and plays a pivotal role in the regulation of the immune response to microbial pathogens [19]. IL-10 is released by monocytes and inhibits production of TNF- α , IL-1 β , and IL-6 [20]. As IL-10 expression is delayed relative to that of TNF- α , IL-1 β , and IL-6, its release provides an efficient autocrine mechanism for controlling proinflammatory cytokine production by monocytes [21]. Although the precise mechanisms by which IL-10 mediates these inhibitory effects have not been determined, both transcriptional and posttranscriptional mechanisms as well as antioxidant effects have been proposed. The balance of pro- and anti-inflammatory cascades may be important determinants of the extent of host inflammatory responses and adverse clinical outcomes. Knowledge gained in the understanding of the role and regulation of IL-10 in acute inflammatory states, for instance, may lead to the development of therapeutic strategies aimed at the modulation of IL-10 synthesis. Of note, the administration of glucocorticoids has been shown to modulate the inflammatory response syndrome in favor of IL-10, resulting in a decrease in circulating TNF- α , IL-6, and IL-8 levels and an increase in IL-10 levels [22, 23].

The IL-1Ra is a member of the IL-1 family that binds to IL-1 receptors but does not transduce intracellular signals [24]. IL-1Ra is mainly secreted by monocytes in response to other cytokines and microbial components. Endogenous IL-1Ra is important in host defense against endotoxin-induced injury as well as in acute and chronic inflammatory states [25]. Studies have shown that a 1000-fold excess of IL-1Ra is required to block the hemody-

namic effects of IL-1 β [26]. Consequently, the balance between proinflammatory cytokines such as TNF- α and IL-1 β , and their specific inhibitors such as IL-10 and IL-1Ra, respectively, may be critical in determining the extent of the inflammatory response, and may have an important influence on outcomes of patients suffering from an acute inflammatory disorder.

Th1 and Th2 responses

Monocyte and T-lymphocyte interactions also play a critical role in the systemic response to acute inflammatory states. Two types of T-helper (Th) cells produce distinct patterns of cytokines. Th1 cells secrete interleukin-2 (IL-2) and interferon-gamma (IFN- γ), whereas Th2 cells mainly produce IL-4 and IL-10. The relative activation of Th cell subsets is determined by a variety of factors, including antigen presentation and the cytokine milieu. In sepsis, the patterns of Th1 and Th2 cell responses are not well known. In one study of healthy volunteers, after low-dose endotoxin administration, the T-cell immune response shifted toward Th2 cell type responses, with depressed circulating IL-2 levels contrasting with marked increases in IL-10 levels [27]. Although the balance of Th1/Th2 cell activation may be important in directing effector pathways and patterns of injury in acute glomerulonephritis [28], the role of lymphocytes in the pathophysiology of acute tubular necrosis is less well understood.

THE INFLAMMATORY NETWORK IN ARF

Sepsis injury in ARF

ARF often develops in the setting of sepsis, and the role of proinflammatory cytokines in the development of endotoxin-mediated ARF is well documented [29–31]. In sepsis, cytokine release is initiated by a number of factors, including bacterial endotoxin, ischemia/reperfusion, complement activation, and the redundant effect of other cytokines and immune modulators (Fig. 3). This results in leukocyte activation along with the expression of adhesion molecules, and the production of various biologically active substances, including oxygen-free radicals, arachidonic acid metabolites, platelet-activating factor, nitric oxide, endothelins, and heat-shock proteins (HSPs) [32, 33]. This proinflammatory cascade contributes to endothelial injury of the renal microvascular bed [34], leading to the development of tubular dysfunction and ARF. In addition, bacterial products can activate neutrophils within an already injured kidney [35], and IL-8 acts to recruit neutrophils to sites of inflammation [36]. Of note, morphologic studies of acute tubular necrosis in humans demonstrate a preponderance of neutrophils in vasa recta and the interstitium, supporting a role for inflammation in a condition that has often been thought to be noninflammatory [37].




Fig. 3. Schematic representation of the inflammatory response to sepsis and resulting kidney injury. (Modified from [205]; reprinted with permission. Jaber BL et al: *Blood Purif* 22: 101–111, 2004). Abbreviations are: GFR, glomerular filtration rate; NAG, N-acetyl- β -D-glucosaminidase; KIM, kidney injury molecule 1.

Ischemic and nephrotoxic injury in ARF

Although systemic or renal hemodynamic insults are important in the pathogenesis of ARF, increasing evidence supports an important role for inflammatory mechanisms in the pathogenesis of ARF following both ischemic [34] and nephrotoxic [38, 39] injury.

In a mice model of whole body ischemia, ARF is characterized by inflammation with increased renal myeloperoxidase (MPO) levels and increased gene expression of intercellular adhesion molecule-1 (ICAM-1) and IL-6 [40]. This inflammatory response as well as the development of ARF is partly reduced in T-cell-deficient mice [40], arguing for the importance of these cells in the pathogenesis of ARF following ischemia/reperfusion injury.

In a mouse model of cisplatin nephrotoxicity, ARF is characterized by the activation of proinflammatory cytokines (TNF- α and IL-1 β mRNA) and chemokines [monocyte chemoattractant peptide-1 (MCP-1) mRNA], and TNF- α urinary levels are increased [39]. The use of pentoxifylline, an inhibitor of TNF- α production, or a TNF- α monoclonal antibody blunts cisplatin-induced increases in TNF- α and IL-1 β mRNA and reduces TNF- α urinary levels. In addition, TNF- α inhibitors ameliorate cisplatin-induced renal dysfunction and reduce structural damage.

Similarly, TNF- α -deficient mice are resistant to cisplatin nephrotoxicity [39]. Overall, these data suggest that TNF- α plays a central role in these responses [39].

Chemokines, including MCP-1 and IL-8, also play a prominent role in the acute inflammatory response in several models of ARF. Indeed, following ischemia/reperfusion injury, kidney MCP-1 mRNA is increased and correlates with the extent of mononuclear infiltrate [41]. Finally, a kidney that is already primed with activated inflammatory cells may be further affected by the immunologic activation induced by the contact of blood with poorly biocompatible dialysis membranes [42].

With a growing body of evidence that there is increased expression of proinflammatory cytokines in experimental models of ARF, the role of anti-inflammatory strategies in limiting tissue injury is emerging. For instance, the exogenous administration of IL-10 inhibits TNF- α and ICAM-1 mRNA expression in mice kidneys following ischemic and nephrotoxic injury [38], and antibody to ICAM-1 protects the kidney against ischemic and nephrotoxic injury [43, 44]. The administration of IL-10 also inhibits macrophage-mediated injury in experimental glomerulonephritis by inhibiting MCP-1, IL-1 β , and ICAM-1 expression [45, 46].




Fig. 4. Key pathways involved in the generation and degradation of oxidants in acute renal failure. Abbreviations are: O_2^- , superoxide anion; SOD, superoxide dismutase; MPO, myeloperoxidase; NO, nitric oxide; $ONOO^-$, peroxynitrite. $OH\cdot$, hydroxyl radical; GPX, glutathione peroxidase; $HOCl$, hypochlorous acid; iNOS, inducible nitric oxide synthase.

In humans, urinary cytokine levels correlate with urinary markers of tubular dysfunction. Indeed, following surgery with cardiopulmonary bypass (CPB), changes in plasma and urinary cytokine homeostasis correlate with urinary N-acetyl- β -D-glucosaminidase (NAG) level, a biomarker of renal proximal tubular dysfunction [47, 48]. Whether urinary cytokines in humans are due to filtration by the glomerulus or production in situ is not clear. They may be involved in the pathobiology of ARF, through paracrine effects. While the role of urinary cytokines and other biomarkers of kidney injury remain unknown, a recent study demonstrated the detection of the soluble form of human kidney injury molecule-1 (KIM-1) in the urine of patients with biopsy-proven acute tubular necrosis [49]. This transmembrane protein is expressed by the human renal proximal tubule, and has been proposed as a useful biomarker for renal proximal tubule injury, to facilitate the early diagnosis of ARF [34, 50].

In humans, plasma cytokine levels have been correlated with the risk of ARF and adverse outcomes. Both plasma IL-6 and IL-10 levels have been shown to differentiate surviving from nonsurviving patients with ARF [51]. Furthermore, among patients with septic shock, elevated serum levels of sTNF-R (forms I and II) were found to be independent predictors for the development of ARF and mortality [52]. In another study of septic patients, an elevated plasma IL-6 to IL-10 ratio was associated with a higher risk of multiple organ failure, suggesting that IL-10 deficiency may contribute to organ dysfunction in sepsis [53].

Oxidative stress in ARF

Under normal physiologic conditions, a homeostatic balance exists between the formation of ROS and their re-

moval by endogenous antioxidant compounds [54], which reside in the intracellular space, the cellular membrane, and the extracellular space.

The reduced form of nicotinamide adenine dinucleotide phosphate (NADPH) oxidases are membrane-associated enzymes that catalyze the production of superoxide and they are highly expressed in neutrophils and endothelial cells. [55]. NADPH oxidase has several subunits, including gp91phox and p22phox, which are electron-transfer proteins. These electron-transfer proteins are predominantly intracellular in endothelial cells, and extracellular in neutrophils [56]. This may explain cell-specific differences in the function of the two enzymes. In sepsis and possibly ARF, there are several potential sources of ROS, including the mitochondrial respiratory electron transport chain, xanthine oxidase activation as a result of ischemia/reperfusion and neutrophil-associated respiratory burst via the activation of membrane-bound NADPH oxidase [57].

MPO is released by activated neutrophils and monocytes and takes part in primary host defenses by catalyzing the production of hypochlorous acid, a potent oxidant (Fig. 4). In addition to its antimicrobial activity, MPO has been involved in a broad range of noninfectious diseases, including ischemia/reperfusion injury, vasculitis, atherosclerosis, cancer, and neurodegenerative disorders [58].

In human cells, glutathione and the glutathione peroxidases (GPX) constitute the principal intracellular antioxidant defense system [59]. There are at least four different glutathione peroxidases, all of which contain selenocysteine at their active sites [60]. Glutathione peroxidase 1 (GPX1), the ubiquitous intracellular form and key antioxidant enzyme within most cells, uses glutathione

(GSH) to reduce hydrogen peroxide to water and lipid peroxides to their respective alcohols [61], and it also reduces peroxynitrite [62]. Whereas GPX1 is particularly abundant in the kidney, GPX2 is expressed primarily in gastrointestinal tissues and GPX3 is extracellular and is present mainly in the plasma. GPX4 is detected in most tissues both in the cytosol and within membranes and has the ability to reduce peroxidized phospholipids in cellular membranes [63].

Superoxide dismutase (SOD) converts superoxide anion to hydrogen peroxide and is represented by three different ubiquitously expressed enzymes: cytosolic copper- and zinc-containing SOD (SOD1), mitochondrial manganese-containing SOD (SOD2), and extracellular SOD (SOD3). Extracellular SOD is most active in the vessel wall and has been shown to regulate the availability of nitric oxide by scavenging superoxide anion [63, 64].

Although β -carotene, lycopene, and coenzyme Q have all been implicated as cell membrane antioxidants, fat-soluble vitamin E or α -tocopherol is by far the most important free radical quencher, residing in the hydrophobic inner layer of membranes [57].

Finally, an additional line of extracellular antioxidant compounds handles the extrusion of ROS into the extracellular space that can occur, for example, following neutrophil activation. Indeed, metal-binding plasma proteins, including lactoferrin and serum albumin, via oxidizable thiol groups, which permit free radical scavenging, are valuable circulating antioxidants. Other important circulating antioxidants include vitamin C, uric acid, and bilirubin [57].

Oxidative stress occurs when a balance is disrupted by excessive production of ROS, including superoxide, hydrogen peroxide, and hydroxyl radicals, and/or by inadequate antioxidant defenses [57], including suboptimal levels of SOD, catalase, GPX, vitamins C and E, and reduced GSH (Fig. 4).

Both of these imbalances may occur in ARF. Indeed, in recent years, considerable evidence has incriminated free oxygen radicals and ROS in the pathogenesis of ischemic and nephrotoxic ARF [65]. Experimental studies indicate that deprivation of antioxidant defenses aggravates renal injury [65]. Strategies to enhance endogenous antioxidant defenses can attenuate renal injury induced by ischemic or nephrotoxic insults [65, 66]. These observations have been supported by data from clinical observational studies indicating depressed antioxidant defenses and increased oxidative stress in critically ill patients with multiple organ failure, including ARF [67–69]. Deficiencies of water-soluble vitamins and trace elements may also occur in patients with ARF undergoing renal replacement therapy [69, 70], which may further compromise antioxidant defenses. Finally, blood-membrane interactions and neutrophil activation during dialysis may lead to the in-

tradialytic release of MPO and generation of ROS, which further exacerbate oxidative stress [69]. This may be an additional source of tissue injury as MPO activity is already enhanced in the renal parenchyma, following ischemia/reperfusion injury, reflecting on the recruitment of activated neutrophils to sites of injury [71].

The use of N-acetyl cysteine (NAC) to prevent ARF has recently emerged. This antioxidant acts primarily by scavenging oxygen free radicals, mainly hydroxyl radicals. This effect is most pronounced in the intracellular space where it also helps restore reduced glutathione stores. In humans, NAC has been shown to be beneficial in preventing contrast-induced nephropathy [72]. In experimental models of ischemic ARF, NAC ameliorates renal vasoconstriction, in part through prostaglandins and nitric oxide [73, 74].

Nitric oxide metabolism in ARF

Nitric oxide is an important signaling molecule that is generated from L-arginine by the action of nitric oxide synthase (NOS), and has a half-life of only a few seconds in vivo. However, it is soluble in both aqueous and lipid media, and therefore, readily diffuses through the cytoplasm and plasma membranes. In the vasculature, nitric oxide stimulates the intracellular production of cyclic guanosine monophosphate (cGMP), resulting in smooth muscle relaxation and vasodilatation. In the extracellular milieu, nitric oxide reacts with oxygen and water to form nitrates and nitrites, two harmless compounds. Nitric oxide toxicity is linked to its ability to combine with superoxide anions to form peroxynitrite, an oxidizing free radical that can cause DNA fragmentation and lipid oxidation. In the mitochondria, peroxynitrite acts on the respiratory chain complex and manganese-containing SOD, to generate superoxide anions and hydrogen peroxide, respectively.

In ischemia/reperfusion injury of the kidney, nitric oxide is produced to a greater degree by tubular and interstitial cells rather than the endothelium. In addition to its role as a strong vasodilator, nitric oxide inhibits production of vasoconstrictors and reduces inflammation by neutralizing ROS [75]. Studies have found that nitric oxide derived from inducible NOS (iNOS) contributes to the preconditioning effect in the kidney [76]. Although nitric oxide is not currently used as a therapy for renal injury [77], there is evidence in humans that the use of L-arginine, the substrate for iNOS-dependent nitric oxide generation, reduces kidney injury [78].

Summary

In summary, although animal and human studies have incriminated cytokines, chemokines, ROS, and nitric oxide in the pathophysiology ARF, the role of susceptibility genetic factors, which may influence the above mentioned

Table 1. Selected list of polymorphism of immune response genes in humans

Gene	Polymorphism			Gene expression or function	Reference
	Site	Class	Position		
Cytokines					
TNF- α	Promoter	SNS (G to A)	−238	Affected	[80]
TNF- α	Promoter	SNS (G to A)	−308	Affected	[82–85]
TNF- β (LT- α)	Intron	SNS (G to A)	+250 (intron 1)	Affected ^a	[87]
TNF- β (LT- α)	Intron	SNS (G to A)	+1069 (intron 1)	Affected ^a	[86]
IL-1 α	Promoter	SNS (C to T)	−889	Affected	[104]
IL-1 α	Intron	46-bpVNTR	Intron 6	Unknown	[206]
IL-1 β	Exon	SNS (C to T)	+3953 (exon 5)	Affected	[207]
IL-1 β	Promoter	SNS (C to T)	−511	Affected	[105]
IL-1Ra	Intron	86-bp VNTR	Intron 2	Affected	[111]
IL-6	Promoter	SNS (G to C)	−174	Affected	[120, 122, 124]
IL-6	Promoter	SNS (G to C)	−572	Affected	[121]
IL-10	Promoter	SNS (G to A)	−1082	Affected	[132, 133]
IL-10	Promoter	SNS (C to T)	−819	Affected	[80, 133]
IL-10	Promoter	SNS (C to A)	−592	Affected	[80]
Chemokines					
IL-8	Promoter	SNS (A to T)	−251	Affected	[140, 141]
IL-8	3'UTR	SNS (G to A)	+2767	Affected	[142, 143]
IL-8	Promoter	SNS (C to T)	−845	Affected	[144]
MCP-1	Promoter	SNS (G to A)	−2518	Affected	[145, 146]
Toll-like receptors (TLR)					
TLR2	Exon	SNS (C to T)	+2029	Affected	[208]
TLR2	Exon	SNS (G to A)	+2251	Affected	[155]
TLR4	Exon	SNS (A to G)	+896	Affected	[152]
Heat shock proteins (HSP)					
HSP70-2	Exon	SNS (G to A)	+1267	Affected	[159]
HSP70-2	Exon	SNS (G to A)	+1538	Unknown	[160]
HSP70-Hom	Exon	SNS (C to T)	+2437	Unknown	[160]
Oxidant stress-related enzymes					
HO-1	Promoter	MSR (GT) _n	−263, −185	Affected	[172, 173]
NADPH oxidase p22phox	Exon	SNS (C to T)	+242 (exon 4)	Affected	[178]
NADPH oxidase p22phox	3'UTR	SNS (A to G)	+640	Unknown	[178]
MPO	Promoter	SNS (G to A)	−463	Affected	[186, 187]
SOD3	Exon	(C to G)	+637	Affected	[188]
Catalase	Promoter	SNS (C to T)	−262	Affected	[189]
GPX1	Exon	MSR (GCG) _n	Exon 1	Unaffected	[190]
GPX1	Exon	SNS (C to T)	+593	Unaffected	[192]
GST M1	Exon ^b	—	—	Affected	[194, 195]
iNOS	Promoter	SNS (C to T)	−1173	Affected	[198]

Abbreviations are: SNS, single nucleotide substitution; VNTR, variable number of tandem repeats; MSR, microsatellite repeats; UTR, untranslated region; HSP, heat shock protein; MPO, myeloperoxidase; SOD, superoxide dismutase; GPX, glutathione peroxidase; GST, glutathione-S transferase; iNOS, inducible nitric oxide synthase; TNF, tumor necrosis factor; IL, interleukin; HO, heme oxygenase; NADPH, nicotinamide adenine dinucleotide phosphate.

^a TNF- α production is affected.

^b Gene deletion prevalent in up to 50% of humans.

specific gene activation pathways, and consequently, both the development and severity of ARF following ischemic or nephrotoxic injury, has remained largely unexplored.

Polymorphisms of selected immune response genes are reviewed in the next section, with special emphasis on those that have been associated with an altered rate of gene expression. Associations between these polymorphisms and acute inflammatory states are summarized, and implications in the field of ARF are hypothesized.

CANDIDATE GENES ALTERING THE INTENSITY OF THE INFLAMMATORY RESPONSE

The study of polymorphism of immune response genes in human disease may enhance the understanding of the cause and pathology of disease, identify potential mark-

ers of susceptibility, severity and clinical outcomes, identify potential markers for responders vs. nonresponders in therapeutic trials, identify targets for therapeutic intervention, and identify novel or improve existing strategies to prevent disease [7, 8].

Host immune responses strongly correlate with the severity and outcome of acute inflammatory and infectious states, and the extent of the inflammatory responses is in part genetically determined. In recent years, the study of polymorphism of immune response genes, including cytokines and other immune modulators, has become the subject of intense interest, as these genetic markers may be potential determinants of susceptibility to or severity of acute illnesses. Polymorphisms of key immune regulatory molecules are reviewed (Tables 1 and 2).

Table 2. Positive associations between polymorphism of immune response genes and acute infectious and inflammatory disorders

Gene	Polymorphic allele	Acute illness	Reference
Cytokines			
TNF- α	–308 A-allele (TNF α 2)	Sepsis/septic shock	[90–92, 97, 98]
TNF- α	–308 A-allele (TNF α 2)	Meningococcal disease	[94]
TNF- α	–308 A-allele (TNF α 2)	Cerebral malaria	[95]
TNF- α	–308 A-allele (TNF α 2)	Mucocutaneous leishmaniasis	[96]
TNF- α	–308 A-allele (TNF α 2)	Rate of body temperature normalization following cardiopulmonary bypass	[102]
TNF- α	–308 A-allele (TNF α 2)	Neonatal acute renal failure	[130]
TNF- α	–308 A-allele (TNF α 2)	Severe acute kidney-pancreas transplant rejection episodes	[135]
TNF- α	–308 A-allele (TNF α 2)	Increased risk of death in dialysis-requiring acute renal failure	[103]
TNF- α	–238 A-allele	Malarial anemia	[209]
TNF- β ^a	+250 AA genotype	Septic shock in community-acquired pneumonia	[93, 163]
TNF- β	+250 G-allele	Prolonged mechanical ventilation following cardiopulmonary bypass	[99]
TNF- β	+1069 (NcO1) allele 2/2	Septic shock following acute biliary pancreatitis	[210]
TNF- β	+1069 (NcO1) allele 2/2	Higher circulating TNF- α levels following cardiopulmonary bypass	[100]
TNF- β	+1069 (NcO1) allele 2/2	Increased mortality in severe sepsis	[97]
TNF- β	+1069 (NcO1) allele 2/2	Susceptibility to severe posttraumatic sepsis	[98]
IL-1 α	–889 TT genotype	Osteomyelitis	[211]
IL-1 β	+3953 TT genotype	Osteomyelitis	[211]
IL-1 β	–511 allele 2/2	Febrile seizure in children	[106]
IL-1 β	+3953 allele 2	Increased risk of ESRD in PR3-ANCA vasculitis	[118]
IL-1 β	–511 allele 1/2	Survival in meningococcal disease	[212]
IL-1Ra	86 bp VNTR (allele 2)	Susceptibility to sepsis	[113, 114]
IL-1Ra	86 bp VNTR (allele 2)	Reduced Mantoux response to purified protein derivative of <i>Mycobacterium tuberculosis</i>	[115]
IL-1Ra	86 bp VNTR (allele 2)	Increased risk of ESRD in PR3-ANCA vasculitis	[118]
IL-1Ra	86 bp VNTR (allele 2)	Severity of Henoch-Schönlein purpura-associated nephritis	[116]
IL-1Ra	86 bp VNTR (allele 2)	High soluble endothelial activation (von Willebrand factor and E-selectin) markers in acute coronary syndromes	[117]
IL-6	–174 GG genotype	Improved survival in sepsis	[125]
IL-6	–174 C-allele	Neonatal acute renal failure	[130]
IL-6	–572 C-allele	Higher serum C-reactive protein level	[121]
IL-10	–1082 GA genotype	Meningococcal disease	[134]
IL-10	–1082 G-allele	Severity of illness in community-acquired pneumonia	[126]
IL-10	–1082 G-allele	Severe acute kidney-pancreas transplant rejection episodes	[135]
IL-10	–1082 GA genotype	Susceptibility to pulmonary tuberculosis	[213]
IL-10	–1082 G-allele	Decreased risk of death in dialysis-requiring acute renal failure	[103]
IL-10	–592 A-allele	Increased mortality in sepsis	[137]
IL-10	–592 AA genotype	Decreased risk of acute GVHD and death	[214]
Chemokines			
IL-8	+2767 A-allele	Henoch-Schönlein purpura-associated nephritis	[143]
IL-8	–251 A-allele	Enterococcal <i>Escherichia coli</i> diarrhea	[140]
IL-8	–251 A-allele	Respiratory syncytial virus bronchiolitis	[141]
IL-8	–845 C-allele	Severity of lupus nephritis	[144]
MCP-1	–2518 G-allele	Premature kidney graft failure	[147]
Toll-like receptors (TLR)			
TLR2	Arg753Gln	Gram-positive bacterial sepsis	[155]
TLR2	Arg677Trp	Susceptibility to lepromatous leprosy	[157]
TLR4	Asp299Gly, Thr399Ile	Gram-negative bacterial sepsis	[154]
TLR4	Asp299Gly	Sepsis and septic shock	[153, 156]
Heat shock proteins (HSP)			
HSP70-2	+1267 GG genotype	Neonatal acute renal failure	[159]
HSP70-2	+1267 AA genotype	Increased risk of septic shock	[163]
HSP70-Hom	+1538 CT genotype	Multiple organ failure following trauma	[160]
Oxidant stress-related enzymes			
HO-1	Long (GT) _n repeats	Increased susceptibility to coronary artery disease in diabetic patients	[173]
HO-1	Long (GT) _n repeats	Increased risk of restenosis following PTCA	[174]
HO-1	Long (GT) _n repeats	Increased risk for developing abdominal aortic aneurysm	[175]
HO-1	Long (GT) _n repeats	Increased susceptibility to pulmonary emphysema in cigarette smokers	[176]
NADPH oxidase p22phox	+242 CC genotype	Increased oxidized high-density lipoprotein cholesterol in type 2 diabetes mellitus	[183]
MPO	–463 GG-genotype	Increased prevalence of cardiovascular disease and higher circulating levels of pentosidine in ESRD	[187]
GST	GST M1-0 genotype	Increased risk for lung cancer in heavy smokers	[196]
GST	GST M1-B allele carrier	Decreased risk of delayed graft function	[197]
iNOS	–1173 T-allele carrier	Decreased risk of malarial complications	[198]

Abbreviations are: ESRD, end-stage renal disease; PR3-ANCA, proteinase 3 antineutrophilic cytoplasmic antibody; PTCA, percutaneous transluminal coronary angioplasty; GVHD, graft versus host disease; GST, glutathione-S transferase; iNOS, inducible nitric oxide synthase; HO, heme oxygenase; MPO, myeloperoxidase; IL, interleukin; TNF, tumor necrosis factor; NADPH, nicotinamide adenine dinucleotide phosphate.

^aTNF- β is lymphotoxin alpha (LT- α).

The TNF locus

The TNF- α and TNF- β [also known as lymphotoxin- α (LT- α)] genes are located on the short arm of chromosome 6. Polymorphism within the 5'-flanking region of the TNF- α gene at positions -238 (G to A) and -308 (G to A) have been reported, and the -308 A-allele, also referred to as the TNF- α 2 allele, has been associated with high promoter activity [79–81]. Moreover, the TNF- α 2 allele has been found to correlate with enhanced spontaneous and stimulated TNF- α production both in vitro [82, 83] and in vivo [84, 85].

The gene coding for TNF- β (LT- α) is located near the TNF- α gene. Single nucleotide polymorphism at position +1069 (G to A) in the first intron of the gene has been characterized [86]. The "A" variant is the TNF- β 2 allele whereas the less frequent "G" variant is known as the TNF- β 1 allele. The TNF- β 1 allele is associated with the presence of a digestion site for the DNA sequence-specific restriction enzyme *Nco*I [87, 88]. This results in a restriction fragment length polymorphism (RFLP). Monocytes from TNF- β 2 homozygous individuals produce significantly higher amount of IL-1 β and TNF- α in response to stimuli [87]. A recent study has demonstrated strong linkage disequilibrium between TNF- α -308 (G to A) promoter and TNF- β *Nco*I polymorphisms, whereby heterozygosity for both TNF polymorphisms was associated with an increased TNF- α response [89]. This finding highlights the importance of linkage disequilibrium in studies examining disease association with polymorphism of individual immune response genes (see below).

Polymorphism in the TNF- α gene has been studied extensively and has been associated with adverse clinical outcomes among patients suffering from sepsis [90–92]. In one such study, the TNF- α high producer genotype (TNF- α 2 allele) was strongly associated with an increased susceptibility to septic shock and mortality [90]. Furthermore, this TNF- α high producer genotype has been associated with increased morbidity and mortality from several acute infectious diseases, including community-acquired pneumonia [93], meningococcal disease [94], malaria [95], and mucocutaneous leishmaniasis [96]. Other studies have linked the TNF- β 2 homozygous carriage, also associated with a TNF- α high producer genotype, to higher fatality among patients with severe sepsis [97], and increased susceptibility to severe post-traumatic sepsis [98].

A few studies have investigated the influence of polymorphisms in the TNF- α [99] and TNF- β [100] genes on the expression of TNF- α in patients undergoing coronary artery bypass graft (CABG) surgery with CPB. Some have also associated the TNF- β +250 G-allele with prolonged mechanical ventilation [99, 101] and the TNF- α -308 G-allele with a slower rate of body temperature normalization following CABG surgery [102]. In one study of patients with dialysis-requiring ARF, carriers of the

TNF- α -308 A-allele had higher TNF- α production by endotoxin-stimulated whole blood leukocytes, a higher APACHE II score and a higher mortality risk [103].

It is clear that these genetically determined differences related to TNF- α gene polymorphisms may influence variations in proinflammatory cytokine responses to stressful stimuli. In patients with ARF, this may have important implications, as the extent of proinflammatory responses may dictate the severity of ARF and requirement for dialysis, as well as overall morbidity and mortality.

The IL-1 gene family

IL-1 is an important mediator of inflammation and tissue damage in multiple organs, both in experimental animal models of disease and in human diseases. The IL-1 gene family consists of IL-1 α , IL-1 β , IL-1 receptors (I and II), and IL-1Ra. The genes encoding the IL-1 α , IL-1 β , and IL-1Ra proteins are located on the short arm of chromosome 2, and the IL-1 receptor proteins on the long arm of chromosome 2.

An analysis of the allelic polymorphism in the promoter region of the IL-1 α gene at position -889 revealed that the TT genotype significantly increased the transcriptional activity of the gene relative to the CC genotype, with a slight increase of the IL-1 α mRNA and plasma protein levels [104].

A SNP has been identified within exon 5 of the IL-1 β gene at position +3953 (C to T), which has been associated with regulation of gene transcription [80]. Another polymorphism has also been described within the 5'-flanking promoter region of the gene at position -511 (C to T) [105]. The IL-1 β -511 polymorphism has been associated with febrile seizures in children [106], variation in CRP level [107], and severity of meningococcal disease [105].

The IL-1Ra gene contains VNTR of 86 bp within intron 2, giving rise to a penta-allelic minisatellite polymorphism. This is in linkage disequilibrium with IL-1Ra polymorphism at position +2018 within exon 2 [108]. The uncommon allele 2, which consists of two 86 bp repeats, has been associated with increased IL-1 β production in vitro [109]. Some, but not all studies have also shown that this allele is associated with increased IL-1Ra production by human monocytes [110, 111]. Of note, a recent study demonstrated that IL-1 β bioactivity is under the genetic control of IL-1Ra [112].

In several studies, the IL-1Ra polymorphic allele-2 has been associated with increased susceptibility to sepsis [113, 114], a reduced response to purified protein derivative of *Mycobacterium tuberculosis* (Mantoux test) [115], and with severity of nephritis in Henoch Schönlein purpura [116]. Among patients with acute coronary syndromes, the IL-1Ra allele-2 has also been associated with higher soluble endothelial activation markers (von

Willebrand factor and E-selectin) and the likelihood of a troponin-positive status compared with noncarriage [117].

Several studies have raised the question of whether polymorphism of IL-1 β and IL-1Ra, known as the IL-1 gene cluster, when examined in conjunction, may have a role in the pathogenesis and/or severity of inflammatory and infectious diseases, by influencing the ratio between these two molecules. This altered ratio may lead to either an attenuated or enhanced inflammatory reaction *in situ* [118].

IL-6

Interleukin-6 is a pleiotropic cytokine with both pro- and anti-inflammatory properties [119]. The IL-6 gene is located on the long arm of chromosome 7. A polymorphism has been identified within the 5'-flanking region of the IL-6 gene at position -174 (G to C) [120] and -572 (G to C) [121]. Transient transfection experiments using constructs containing either allele of the -174 5'-flanking region polymorphism indicate that the C allele results in suppression of IL-6 transcription in response to endotoxin or IL-1 β [120]. The presence of the C allele has also been associated with lower serum levels of IL-6 in healthy subjects supporting the *in vitro* observation of reduced promoter strength [120]. In contrast, carriers of the wild-type IL-6 -174 G-allele have higher serum IL-6 levels [122].

The frequency of the IL-6 -174 C-allele has been estimated at 40% in an Anglo-Saxon Caucasian population [120]. Limited data suggest, for instance, that ethnic differences exist in regard to IL-6 genotype between Caucasian and African American populations in the United States [123].

Studies linking the IL-6 -174 promoter region polymorphism to clinical outcomes in acute inflammatory states have yielded conflicting results. In severely injured blunt trauma patients, there was no association between the IL-6 -174 promoter genotypes and *ex vivo* whole blood IL-6 production levels after stimulation with endotoxin [124]. In other studies, this polymorphism was not associated with increased incidence of sepsis nor did it correlate with plasma IL-6 level [125]. This polymorphism also failed to predict severity of illness in community-acquired pneumonia [126] and episodes of acute renal allograft rejection [127].

These studies contrast with a few reports demonstrating an association between this polymorphism and acute inflammatory disorders. In one study, there was a significant association between the IL-6 -174 GG homozygous genotype and improved survival in sepsis [125]. This was independent of the systemic IL-6 response, arguing for the role of other genetically linked polymorphisms that might have been the underlying cause for this association.

The IL-6 -572 C-allele and -174 CC genotypes have correlated with perioperative IL-6 profiles among patients undergoing CABG surgery [128]. This contrasts with one study where among patients undergoing CABG surgery, the IL-6 -174 GG homozygous genotype was associated with higher acute phase levels of IL-6 and with longer stays in the hospital and in the intensive care unit compared with patients carrying the -174 C-allele [129].

The constellation of high TNF- α producer and low IL-6 producer genetic variants, which may lead to higher pro- and lower anti-inflammatory capacities, has been associated with an increased risk for ARF in low-birth-weight infants [130]. This is not inconceivable as high-dose IL-6 stimulates tubular regeneration in rat kidneys following nephrotoxic injury [131], thus providing an additional effect by which IL-6 may influence the course of ARF.

In summary, the variability in association between IL-6 gene polymorphism and acute inflammatory disorders may stem from its known pro- and anti-inflammatory properties and the high frequency of the wild-type in the general population (i.e., the high IL-6 production phenotype).

IL-10

The IL-10 gene is located on the long arm of chromosome 1. Studies have suggested that 70–80% of the variation in IL-10 production is genetically determined and appears to be controlled at the transcriptional level [132]. The IL-10 5'-flanking region, which controls transcription, is polymorphic with a single base pair substitution at position -1082 (G to A) [133]. *In vitro* studies have identified three phenotypic secretion levels for IL-10 based on allelic substitutions (G to A) at the -1082 position: high or intermediate producers (-1082 G-allele carrier, or genotypes GG for high and GA for intermediate producers), and low producers (genotype AA) [80, 133]. These findings have been confirmed by transient transfection studies of monocytic cell lines using constructs of each haplotype [133]. Additional polymorphisms at positions -819 (C to T) and -592 (C to A) have been described but have variably been linked to regulation of gene transcription [80, 133]. Polymorphisms at these two sites are in linkage disequilibrium.

IL-10 -1082 polymorphic alleles have been associated with susceptibility to meningococcal disease and adverse clinical outcomes [134], and severity of illness in community-acquired pneumonia [126]. An increased incidence of acute kidney-pancreas transplant rejection has been observed among organ recipients, as the IL-10 production phenotype increases (low, intermediate, high), but only among those with the TNF- α low producer phenotype [135]. The role of IL-10 as a costimulatory agent for antigen-presenting cells is well known [136]. Such data support the hypothesis that the extent of the alloimmune

response following organ transplantation is in part genetically determined, and that the IL-10 high producer phenotype in this particular setting may be deleterious.

The IL-10 -592 A-allele has been associated with higher mortality in sepsis [137]. In one study of patients with dialysis-requiring ARF, the IL-10 -1082 G-allele carriage was associated with higher IL-10 production and a lower risk of death after adjustment for APACHE II score, multiple organ failure score, and sepsis [103].

Chemokines

IL-8 and MCP-1 are two potent neutrophil and monocyte chemokines, respectively, which have been implicated in the pathogenesis of inflammation in human glomerulonephritis and acute renal allograft rejection [138]. Several studies have attempted to link interpatient variations in the extent of kidney injury to chemokine gene polymorphism, as certain polymorphic alleles may influence the level of chemokine expression in situ, thereby affecting the severity of renal injury.

The IL-8 and MCP-1 genes are located on the short arm of chromosome 4 and 17, respectively [139]. Polymorphism in the IL-8 promoter region at position -251 has been associated with both enteroaggregative *Escherichia coli* diarrhea [140] and respiratory syncytial virus bronchiolitis [141]. A polymorphism at position +2767 (G to A) in the 3'-UTR region of the IL-8 gene has been shown to contribute to its posttranscriptional regulation [142], and an increased frequency of the +2767 A-allele has been found among patients with Henoch Schönlein purpura who develop nephritis [143]. Finally, a novel polymorphism in the promoter region of the IL-8 gene at position -845, found primarily in African Americans, has been associated with the severity of lupus nephritis [144].

A genetic polymorphism in the 5'-flanking region of the MCP-1 gene at position -2518 has been shown to affect MCP-1 expression [145]. Indeed, the MCP-1 -2518 G-allele was associated with higher MCP-1 production levels by human monocytes [145]. Furthermore, the MCP-1 -2518 GG genotype has been associated with higher plasma and urinary MCP-1 levels in patients with lupus nephritis [146], and has been linked to a higher risk for premature kidney graft failure [147].

Toll-like receptors

Toll-like receptors (TLR) are transmembrane proteins that are key mediators of the innate immune response to microbial pathogens. They are responsible for the recognition and initiation of cell-mediated immune responses, such as the release of inflammatory cytokines, phagocytosis, and the expression of costimulatory molecules needed by antigen-presenting cells. Ten different subtypes of TLRs (TLR 1 to 10) have been so far identified in mammals, and these are responsible for the recognition

of separate classes of pathogens that have common specific molecular features, known as pathogen-associated molecular patterns (PAMP).

TLR2 and TLR4 have received particular attention because of their specificity for the recognition of peptidoglycans, lipoteichoic acid, and other antigens specific for gram-positive bacteria, and lipopolysaccharide (LPS) and other bacterial glycolipids specific for gram-negative bacteria [148], respectively. Genetic defects of TLR4 receptors in inbred mouse strains cause resistance to LPS challenge [149]. This lack of an inflammatory response to LPS has been reproduced with TLR4 knockout mice [150]. On the other hand, overexpression of TLR4 in transgenic mice results in an overwhelming inflammatory response after LPS exposure [151].

Polymorphisms of both the TLR2 and TLR4 gene, affecting bacterial antigen responsiveness, have recently been identified in humans (Tables 1 and 2) [152]. These polymorphic alleles have been linked to adverse clinical outcomes in sepsis [153, 154], gram-positive [155] and gram-negative [156] bacterial infections, as well as in mycobacterial infection [157].

Heat shock proteins

HSPs are highly conserved proteins that are expressed in both physiologic and pathologic conditions. HSPs are classified on the basis of molecular weight and the 70 kD proteins referred as the HSP70 family represent the most prominent group, and include constitutive HSP73 and the stress inducible HSP72, HSP70-2, and HSP70-Hom [158–160]. Members of the HSP70 family have been incriminated in the protection against cellular damage from stressful stimuli by acting as chaperone molecules, binding to and preventing aggregation of denatured or abnormal proteins, and facilitating the restoration of normal protein function. Recent studies have shown that HSP70 proteins also act like cytokines and stimulate monocytes to produce proinflammatory cytokines, including TNF- α , IL-1 β , and IL-6 [161]. The increased expression of HSP attenuates experimental renal ischemic injury, and therefore, its modulation by pharmacologic maneuvers may be critical in inflammatory conditions and organs prior to a potential acute insult [162].

The genes encoding for HSP70 proteins are located on chromosome 6. A SNP has been identified within the exon region of the HSP70-2 gene at position +1267 (G to A) [159] and +1538 (G to A) [160]. Another polymorphism has also been described within the codon region of the HSP70-Hom gene at position +2437 (C to T) [160]. Neonates carrying the HSP70-2 (+1267) GG genotype, which is associated with low inducibility of HSP70-2, are at greater risk for developing ARF [159]. This contrasts with a study of adults with community-acquired pneumonia where the HSP70-2 (+1267) AA genotype was

associated with an increased risk of septic shock [163]. Finally, the HSP70-Hom (+1538) CT genotype has been associated with multiple organ failure following trauma [160].

Heme oxygenase-1

Heme oxygenase (HO) is the rate-limiting enzyme in heme catabolism, which leads to the generation of biliverdin, free iron, and carbon monoxide. Three mammalian HO isoforms have been identified, one of which, HO-1, is a stress-responsive protein induced by various oxidative agents. The kidney is one of the prominent sites for intense oxidative processes in the body and is, therefore, extremely vulnerable to free radical-mediated injury [164]. Studies over the past decade indicate that HO-1 is induced as a beneficial response in cells exposed to a diverse array of toxic insults [165]. HO-1 overexpression using gene transfer protects rat livers as well as rat kidney transplants from ischemia/reperfusion injury [166]. Increased HO-1 expression is also cytoprotective in heme-mediated (e.g., rhabdomyolysis) as well as nonheme-mediated (e.g., cisplatin-induced injury) models of ARF [165]. In addition, analyses of HO-1-null mice as well as the first reported HO-1-deficient human have also emphasized the potent anti-inflammatory properties of HO-1. Both mice and humans deficient in HO-1 expression have a phenotype of an increased inflammatory state [167, 168]. Recent studies have focused on the mechanisms by which HO-1 mediates its anti-inflammatory effects. Carbon monoxide, one of the three byproducts of heme catabolism by HO, mediates potent anti-inflammatory effects through a mitogen-activated protein kinase-dependent pathway [169]. Further, other studies have shown that IL-10 induces expression of HO-1 via a p38 mitogen-activated protein kinase-dependent pathway [170]. The inhibition of HO-1 protein synthesis or activity significantly reverses the inhibitory effect of IL-10 on LPS-induced TNF- α production [170]. In addition, IL-10-mediated protection against LPS-induced septic shock is significantly attenuated by cotreatment with the HO inhibitor, zinc protoporphyrin [170]. These observations indicate that HO-1 is an important downstream effector of IL-10. The human HO-1 gene has been mapped to the long arm of chromosome 22 [171] and a highly length polymorphic (GT)_n dinucleotide repeat has been identified in the proximal promoter region [172]. Transient transfection assays with HO-1 promoter/luciferase fusion constructs carrying various lengths of (GT)_n repeats indicate that longer (GT)_n repeats exhibit lower transcriptional activity [173]. Patients with type 2 diabetes mellitus carrying longer (GT)_n repeats have been shown to exhibit higher oxidative stress and increased susceptibility to the development of coronary artery disease [173]. Long (GT)_n repeats have also

been associated with increased risk of restenosis after percutaneous transluminal angioplasty [174], development of abdominal aortic aneurysm [175], and increased susceptibility to pulmonary emphysema in cigarette smokers [176]. The role of this HO-1 promoter (GT)_n polymorphism in modulating susceptibility to or severity of ARF has not been investigated. However, given the cytoprotective effects of HO-1 in experimental models of ARF, it could potentially be an important genetic factor influencing the pathobiology of acute renal injury.

Oxidant stress-related enzymes

Oxygen free radicals are important mediators of cellular injury following organ ischemia and reperfusion [177]. Although polymorphisms affecting key pro- and anti-oxidant enzymes may alter the susceptibility to oxidative stress-mediated injury, the use of genetic epidemiology for the study of oxidative stress-related genes has received little attention in the setting of an acute illness [63]. We will review a selected number of polymorphisms of key oxidant stress-related genes, which may be functionally relevant in human disease.

Given its importance in generating superoxide during the respiratory burst of neutrophils, one might anticipate that polymorphisms disrupting the function of NADPH oxidase might alter oxidative stress-mediated injury in acute inflammatory disorders. Polymorphisms in the gene encoding the NADPH oxidase p22phox subunit have been described. Among them, a single nucleotide substitution (C to T) within exon 4 at position +242, resulting in an amino acid substitution at position 72 (histidine to tyrosine), has been shown to modulate enzyme activity through heme-binding affinity [178]. In one study, the p22phox +242 CC genotype was associated with higher basal and stimulated superoxide generation by human endothelial cells, compared with the p22phox +242 T allele [179]. Another single nucleotide substitution (A to G) at position +640 located in the 3'-UTR of the p22phox gene has also been described, but is of unclear clinical significance [178]. More recent studies have not supported a functional role for the A640G or C242T polymorphisms either in determining endothelial function in patients with coronary artery disease [180] or hypercholesterolemia [181]. Studies on the influence of the p22phox C242T polymorphism on circulating markers of lipid peroxidation [i.e., malondialdehyde (MDA) levels] have yielded conflicting results among patients with coronary artery disease and type 2 diabetes mellitus [182, 183].

With respect to kidney disease, the NADPH oxidase p22phox subunit is expressed by human glomerular mesangial cells, and through the generation of superoxide, may contribute to glomerular injury. Unfortunately, the p22phox C242T polymorphism has not been associated with the severity of glomerulonephritis [184]

or with the development of pre-eclampsia [185]. Despite these negative studies, the neutrophil-associated NADPH oxidase system may be extremely important in sepsis, whereby the respiratory burst during ischemia/reperfusion injury may be in part genetically predetermined. To date, no studies have attempted to link the p22phox C242T polymorphism with the susceptibility to or severity of acute organ injury in sepsis.

MPO is another neutrophil-associated enzyme, which is an important mediator of cellular injury through the generation of oxidants. Variations in the gene encoding for MPO may influence protein expression, and therefore, may be linked to pathologic states through modulation of oxidative activity. There is a well-described SNP within the promoter region of the MPO gene at position -463, consisting of a G to A substitution. The G allele (in contrast to A allele) is associated with high transcriptional activity [186]. In a recent study of patients with chronic kidney failure, carriers of the MPO -463 GG-genotype (associated with higher MPO levels) had a higher prevalence of cardiovascular disease and higher circulating levels of pentosidine, a surrogate marker of protein oxidation [187]. It remains to be seen whether polymorphism of this critical neutrophil-associated enzyme is important in dictating severity of injury following ischemia/reperfusion in ARF.

The SOD3 gene is located on the long arm of chromosome 4 and contains 3 exons. A single nucleotide substitution (C to G) within exon 3 at position +637, resulting in an amino acid substitution at position 213 (arginine to glycine), has been associated with enhanced plasma levels of extracellular SOD [188].

The catalase gene is located on the short arm of chromosome 11 and contains 13 exons. A functional polymorphism in the human catalase promoter, consisting of a C to T substitution at position -262, confers increased basal expression in different cell types and higher enzyme activity in red blood cells [189].

The GPX1 gene is located on the short arm of chromosome 3 and contains 2 exons. A microsatellite polymorphism involving a variable number of three nucleotides repeats (GCG)_n (poly-alanine) in exon 1 has been described [190]. Three polymorphic alleles consisting of 5, 6, and 7 alanine repeats, however, have not been associated with altered GPX1 activity [191]. Another polymorphism, consisting of a C to T substitution at position +593, resulting in an amino acid change at position +197 (proline to leucine) has also been described, but is not associated with altered GPX1 activity [192].

Glutathione S transferases (GSTs) are soluble enzymes that play an important role in the detoxification of oxidative stress end-products including lipid and DNA hydroperoxides. There are at least four GST isoenzymes (M, P, T, and Z) with many described polymorphic alleles [63]. The gene coding for GST M1, the most studied

compound, is located on the short arm of chromosome 1. Three GST M1 alleles have been identified, GST M1-0, GST M1-A, and GST M1-B [193]. Fifty percent of humans lack this enzyme (GST M1-0) [194], whereas the GST M1-A and GST M1-B alleles differ at amino acid residue 172, where lysine is replaced by asparagine, a substitution that has no impact on the enzymatic activity [195]. Smokers who are GST M1 deficient (GST M1-0) are at higher risk of developing lung cancer [196]. In one study of genetic determinants of delayed graft function following kidney transplantation, patients receiving a kidney from a donor who expressed GST M1-B either alone or in combination with GST M1-A experienced significantly lower rates of delayed graft function [197]. These promising data suggest that this antioxidant gene may be protective against ischemia/reperfusion injury following organ transplantation, and may also be applicable to ischemic and nephrotoxic ARF.

Nitric oxide synthase

In the kidney and the vasculature, nitric oxide is produced by three major NOS isoforms: neuronal, endothelial, and inducible NOS (nNOS, eNOS and iNOS, respectively). Neuronal NOS and eNOS participate in the homeostatic control of the vascular tone, the glomerular microcirculation and other processes. In contrast, iNOS is typically induced by proinflammatory stimuli to produce large amounts of nitric oxide, which contributes to the hemodynamic collapse of sepsis and inflammatory injury responses of the glomerulus, vasculature, and the tubulointerstitium.

A single nucleotide substitution (C to T) in the promoter region of the iNOS isoform-2 (iNOS2) gene at position -1173 has been shown to affect nitric oxide expression [198]. Indeed, the iNOS -1173 T-allele carrier has been associated with increased fasting urinary and plasma nitric oxide metabolite concentrations and with a lower risk of malarial complications [198]. To date, there are no known associations between iNOS polymorphism and the development or severity of ARF.

LIMITATIONS OF GENETIC STUDIES

Polymorphism-association studies have traditionally consisted of case-control studies, comparing the prevalence of a genetic marker in unrelated people with a given disease to the prevalence in a control healthy population. In general, such studies have been constrained by small sample sizes, with chance accounting for the results, and therefore, they must be interpreted with caution, especially when biologic plausibility has not been determined or is unknown. A better understanding of genetically linked polymorphisms may be critically important, as the polymorphism of a particular gene may

just be a marker for another, yet to be identified, disease-causing sequence variant [199]. This more comprehensive approach requires the use of linkage analyses and a good understanding of neighboring candidate genes located on the same chromosome. One such good example is the cluster of the IL-1 gene family, with several genes located in proximity on a single chromosome. Furthermore, most published studies describing the frequency of cytokine gene polymorphisms in the general population are biased due to a lack of consideration of ethnicity and geographic boundary [200]. Indeed, certain genetic variants are infrequent in some but not other ethnic groups.

The HapMap Project is an international initiative, with the long-term goal of determining the common patterns of DNA sequence variation in the human genome. This ambitious program plans to determine the genotypes of one million or more sequence variants and their frequencies in DNA samples from populations with ancestry from parts of Africa, Asia, and Europe [201]. Through these population-based genetic studies, the HapMap will allow the discovery of sequence variants that affect common disease across ethnicity and geographic boundary. This endeavor will also facilitate the development of diagnostic tools and targeted therapies.

Since genetic markers are present at birth and predate onset of disease, confounding factors play a minor role unless there is a strong interaction between these markers and an important acquired risk factor [200]. Finally, survival bias is a real concern when genetic markers are being evaluated as predictors of mortality. To circumvent this important limitation, this would require the use of case control studies with matching for age and follow-up time, or more ideally, analytic prospective cohort studies should be employed in an attempt to link polymorphic genes to clinical outcomes over time [202].

Rosenthal and Schwartz [203] have proposed four criteria that need to be fulfilled in establishing medically useful links between genetic polymorphism and human disease. These authors stressed that the genetic variant under study must cause a relevant alteration in the function or level of the gene product. The number of cases associating an allele with a particular phenotype must be large enough to be convincing. The beneficial and harmful phenotypes under study must have clear-cut clinical differences and unequivocal allelic group assignment. Finally, the plausibility of the hypothesis must be convincing.

An important criterion that has also been proposed for the study of genetic polymorphisms is the reproduction of the results in different populations. Other issues to consider when analyzing the results of SNP-association studies pertain to the population characteristics, including the choice of a control group (in case of case-control studies), the prospective or retrospective design of the study, the size, and admixture of the study population,

Table 3. The future of genetic studies in acute renal failure

Large prospective cohort studies
Careful phenotypic characterization
Analysis of genetic patterns and correlation with phenotypic patterns
Inclusion of genetic risk into an acute renal failure-risk algorithm
Use in risk quantification
Use in targeting of therapeutic strategies
“GeneChip” studies

the precise phenotypic characterization, and the effects of confounders. Finally, the demonstration of the strength of an association, the specificity of the effect, the presence of a biologic gradient and gene-dose effect and the notion of temporality all need to be considered.

POTENTIAL RELEVANCE OF GENE POLYMORPHISM IN ARF

ARF is influenced by acquired risk factors such as an exposure to a nephrotoxic drug, and arguably by genetic risk factors that are present at birth. The pathobiology of ARF suggests that products of many genes act concomitantly, culminating in a beneficial or deleterious balance of pro- and anti-inflammatory molecules, which in turn, determines the extent of tissue injury. The number of recognized gene polymorphisms is growing daily. Therefore, future studies should focus on the investigation of combinations of genetic markers. Some have proposed that compilation of “genetic patterns” could be associated with different “clinical patterns” [88], such as for instance, an increased susceptibility or resistance to ARF.

The future of genetic studies in ARF is summarized in Table 3. In brief, genetic epidemiology studies may help unravel the relative importance of genetic markers in predicting the development of hospital-acquired ARF. This would require large prospective cohort studies with strong correlation between these genetic markers and the early detection of urinary biomarkers of kidney injury, a rise in serum creatinine level and dialysis requirement. In the event of successful associations between these markers and outcomes, this would provide the basis for additional studies where this genetic information could be used to stratify patients who are at the highest risk of developing ARF. More important, this knowledge may have far reaching consequences and help identify target patients in clinical trials undergoing high-risk procedures who may benefit from immunomodulatory therapeutic strategies, on the basis of their genotype profiles, with the hope of preventing or attenuating kidney injury.

CONCLUSION

In summary, in vitro and animal studies have incriminated cytokines and other immunoregulatory molecules as important mediators of kidney injury. The relative

importance of genetic factors such as polymorphisms involving immune response genes, in influencing susceptibility to ARF and adverse clinical outcomes, has remained largely unexplored and merits full attention.

Based on the strength of associations observed in studies between gene polymorphisms and the susceptibility to or severity of acute illness and adverse outcomes, we believe that some promising and emerging candidate genes to study in ARF would include the TNF and IL-10 gene [103, 204].

ACKNOWLEDGMENT

Dr. Bertrand L. Jaber is supported by National Institutes of Health Grant K23 DK065102-01.

Reprint requests to Bertrand L. Jaber, M.D., Division of Nephrology, Caritas St. Elizabeth's Medical Center, 736 Cambridge Street, Boston, MA 02135.

E-mail: Bertrand.jaber@cchcs.org

REFERENCES

- NASH K, HAFEEZ A, HOU S: Hospital-acquired renal insufficiency. *Am J Kidney Dis* 39:930–936, 2002
- NOLAN CR, ANDERSON RJ: Hospital-acquired acute renal failure. *J Am Soc Nephrol* 9:710–718, 1998
- LEVY EM, VISCOLI CM, HORWITZ RI: The effect of acute renal failure on mortality: A cohort analysis. *JAMA* 275:1489–1494, 1996
- CHERTOW GM, LEVY EM, HAMMERMEISTER KE, et al: Independent association between acute renal failure and mortality following cardiac surgery. *Am J Med* 104:343–348, 1998
- VENTER J, ADAMS M, MYERS E: The sequence of the human genome. *Science* 291:1304–1351, 2001
- GUTTMACHER A, COLLINS F: Genomic medicine: Genomic medicine—A primer. *N Engl J Med* 347:1512–1520, 2002
- BIDWELL J, KEEN L, GALLAGHER G, et al: Cytokine gene polymorphism in human disease: On-line databases, supplement 1. *Genes Immunol* 2:61–70, 2001
- HAUKIM N, BIDWELL J, SMITH A, et al: Cytokine gene polymorphism in human disease: On-line databases, supplement 2. *Genes Immunol* 3:313–330, 2002
- GENETIC VARIATION IN INDIVIDUALS: Mutation and Polymorphism, in *Thompson & Thompson Genetics in Medicine* (6th ed), edited by Nussbaum R, McInnes R, Willard H, Sixth ed, Philadelphia, WB Saunders Company, 2001, pp 79–94
- VAN DEVENTER S: Cytokine and cytokine receptor polymorphisms in infectious disease. *Intensive Care Med* 26(Suppl 1):S98–S102, 2000
- AMERICAN COLLEGE OF CHEST PHYSICIANS/SOCIETY OF CRITICAL CARE MEDICINE CONSENSUS CONFERENCE: Definitions of sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. *Crit Care Med* 20:864–874, 1992
- JABER BL, PEREIRA BJG: Inflammatory mediators in sepsis: Rationale for extracorporeal therapies. *Am J Kidney Dis* 28:S35–S49, 1996
- MARTIN GS, MANNINO DM, EATON S, MOSS M: The epidemiology of sepsis in the United States from 1979 through 2000. *N Engl J Med* 348:1546–1554, 2003
- GABAY C, KUSHNER I: Acute-phase proteins and other systemic responses to inflammation. *N Engl J Med* 340:448–454, 1999
- KOCH T: Origin and mediators involved in sepsis and the systemic inflammatory response syndrome. *Kidney Int* 53 (Suppl 64):S66–S69, 1998
- PINSKY MR, VINCENT JL, DEVIERE J, et al: Serum cytokine levels in human septic shock. Relation to multiple-system organ failure and mortality. *Chest* 103:565–575, 1993
- CASEY LC BR, BONE RC: Plasma cytokine and endotoxin levels correlate with survival in patients with the sepsis syndrome. *Ann Intern Med* 119:771–778, 1993
- OPAL S, DEPALO V: Anti-inflammatory cytokines. *Chest* 117:1162–1172, 2000
- OPAL SM, HUBER C: The role of interleukin-10 in critical illness. *Curr Opin Infect Dis* 13:221–226, 2000
- VAN DER POLL T, JANSEN J, LEVI M, et al: Regulation of interleukin 10 release by tumor necrosis factor in humans and chimpanzees. *J Exp Med* 180:1985–1988, 1994
- DONNELLY RP, FREEMAN SL, HAYES MP: Inhibition of IL-10 expression by IFN- γ up-regulates transcription of TNF- α in human monocytes. *J Immunol* 155:1420–1427, 1995
- TABARDEL Y, DUCHATEAU J, SCHMARTZ D, et al: Corticosteroids increase blood interleukin-10 levels during cardiopulmonary bypass in men. *Surgery* 119:76–80, 1996
- FILLINGER MP, RASSIAS AJ, GUYRE PM, et al: Glucocorticoid effects on the inflammatory and clinical responses to cardiac surgery. *J Cardiothorac Vasc Anesth* 16:163–169, 2002
- AREND WP, SMITH MFJ, JANSON RW, JOSLIN FG: IL-1 receptor antagonist and IL-1b product in human monocytes are regulated differently. *J Immunol* 147:1530–1536, 1991
- AREND W, MALYAK M, GUTHRIDGE C, GABAY C: Interleukin-1 receptor antagonist: Role in biology. *Annu Rev Immunol* 16:27–55, 1998
- PEREIRA BJ: Cytokine production in patients on dialysis. *Blood Purif* 13:135–146, 1995
- ZIMMER S, POLLARD V, MARSHALL G, et al: Effects of endotoxin on the Th1/Th2 response in humans. *J Burn Care Rehab* 17:491–496, 1996
- TIPPING P, HOLDSWORTH S: T cells in glomerulonephritis. *Springer Semin Immunopathol* 24:377–393, 2003
- WIECEK A, ZEIER M, RITZ E: Role of infection in the genesis of acute renal failure. *Nephrol Dial Transplant* 9:40–44, 1994
- SANTOS OFP, BOIM MA, SCHOR N: Acute renal failure and sepsis—Pathophysiology. *Kidney: A Current Survey of World Literature* 4:59–62, 1995
- CUNNINGHAM P, DYANOV H, PARK P, et al: Acute renal failure in endotoxemia is caused by TNF acting directly on TNF receptor-1 in kidney. *J Immunol* 168:5817–5823, 2002
- FLO TH, HALAAS O, TORP S, et al: Differential expression of toll-like receptor 2 in human cells. *J Leukoc Biol* 69:474–481, 2001
- DYBDAHL B, WAHBA A, LIEN E, et al: Inflammatory response after open heart surgery: Release of heat-shock protein 70 and signaling through toll-like receptor-4. *Circulation* 105:685–690, 2002
- BONVENTRE J, WEINBERG J: Recent advances in the pathophysiology of ischemic acute renal failure. *J Am Soc Nephrol* 14:2199–2210, 2003
- LINAS SL, WHITTEBURG D, PARSONS PE, et al: Mild renal ischemia activates primed neutrophils to cause acute renal failure. *Kidney Int* 42:610–616, 1992
- ZOJA C, ANGIOLETTI S, DONADELLI R, et al: Shiga toxin-2 triggers endothelial leukocyte adhesion and transmigration via NF-kappaB dependent up-regulation of IL-8 and MCP-1. *Kidney Int* 62:846–856, 2002
- SOLEZ K, MOREL-MAROGER L, SRAER J-D: The morphology of acute tubular necrosis in man: Analysis of 57 renal biopsies and a comparison with glycerol model. *Medicine* 58:362–376, 1979
- DENG J, KOHDA Y, CHIAO H, et al: Interleukin-10 inhibits ischemic and cisplatin-induced acute renal injury. *Kidney Int* 60:2118–2128, 2001
- RAMESH G, REEVES W: TNF-alpha mediates chemokine and cytokine expression and renal injury in cisplatin nephrotoxicity. *J Clin Invest* 110:835–842, 2002
- BURNE-TANEY M, KOFLER J, YOKOTA N, et al: Acute renal failure after whole body ischemia is characterized by inflammation and T cell-mediated injury. *Am J Physiol Renal Physiol* 285:F87–F94, 2003
- SUNG F, ZHU T, AU-YEUNG K, SIOW YOK: Enhanced MCP-1 expression during ischemia/reperfusion injury is mediated by oxidative stress and NF-kappaB. *Kidney Int* 62:1160–1170, 2002
- KARSOU S, JABER BL, PEREIRA BJ: Impact of intermittent hemodialysis variables on clinical outcomes in acute renal failure: A critical review. *Am J Kidney Dis* 35:980–991, 2000

43. KELLY KJ, WILLIAMS WWJ, COLVIN RB, BONVENTRE JV: Antibody to inter-cellular adhesion molecule 1 protects the kidney against ischemic injury. *Proc Natl Acad Sci USA* 91:812–816, 1994
44. KELLY K, MEEHAN S, COLVIN R, et al: Protection from toxicant-mediated renal injury in the rat with anti-CD54 antibody. *Kidney Int* 56:922–931, 1999
45. HUANG X, KITCHING A, TIPPING P, HOLDSWORTH S: Interleukin-10 inhibits macrophage-induced glomerular injury. *J Am Soc Nephrol* 11:262–269, 2000
46. KITCHING A, KATERELOS M, MUDGE S, et al: Interleukin-10 inhibits experimental mesangial proliferative glomerulonephritis. *Clin Exp Immunol* 128:36–43, 2002
47. GORMLEY SM, MCBRIDE WT, ARMSTRONG MA, et al: Plasma and urinary cytokine homeostasis and renal dysfunction during cardiac surgery. *Anesthesiology* 93:1210–1216, 2000
48. GORMLEY SM, MCBRIDE WT, ARMSTRONG MA, et al: Plasma and urinary cytokine homeostasis and renal function during cardiac surgery without cardiopulmonary bypass. *Cytokine* 17:61–65, 2002
49. HAN WK, BAILLY V, ABICHANDANI R, et al: Kidney Injury Molecule-1 (KIM-1): A novel biomarker for human renal proximal tubule injury. *Kidney Int* 62:237–244, 2002
50. ICHIMURA T, HUNG C, YANG S, et al: Kidney injury molecule-1 (Kim-1): A tissue and urinary biomarker for nephrotoxicant-induced renal injury. *Am J Physiol Renal Physiol* 286:F552–F563, 2004
51. SIMMONS E, HIMMELFARB J, SEZER M, et al: Plasma cytokine levels predict mortality in patients with acute renal failure. *Kidney Int* 65:1357–1365, 2004
52. IGLESIAS J, MARIK PE, LEVINE JS, NORASEPT II STUDY INVESTIGATORS: Elevated serum levels of the type I and type II receptors for tumor necrosis factor- α as predictive factors for ARF in patients with septic shock. *Am J Kidney Dis* 41:62–75, 2003
53. LOISA P, RINNE T, LAINE S, et al: Anti-inflammatory cytokine response and the development of multiple organ failure in severe sepsis. *Acta Anaesthesiol Scand* 47:319–325, 2000
54. GUTTERIDGE J, MITCHELL J: Redox imbalance in the critically ill. *Br Med Bull* 55:49–75, 1999
55. AZUMI H, INOUE N, TAKESHITA S, et al: Expression of NADH/NADPH oxidase p22phox in human coronary arteries. *Circulation* 100:1494–1498, 1999
56. WHITEHEAD A, FITZGERALD G: Twenty-first century phox: Not yet ready for widespread screening. *Circulation* 103:7–9, 2001
57. MACDONALD J, GALLEY H, WEBSTER N: Oxidative stress and gene expression in sepsis. *Br J Anaesth* 90:221–232, 2003
58. HOY A, LEININGER-MULLER B, KUTTER D, et al: Growing significance of myeloperoxidase in non-infectious diseases. *Clin Chem Lab Med* 40:2–8, 2002
59. URSINI F, MAIORINO M, BRIGELIUS-FLOHE R, et al: Diversity of glutathione peroxidases. *Methods Enzymol* 252:38–53, 1995
60. ARTHUR J: The glutathione peroxidases. *Cell Mol Life Sci* 57:1825–1835, 2000
61. FLOHE L: Glutathione peroxidase. *Basic Life Sci* 49:663–668, 1988
62. SIES H: Glutathione and its role in cellular functions. *Free Radic Biol Med* 27:916–921, 1999
63. FORSBERG L, DE FAIRE U, MORGENTERN R: Oxidative stress, human genetic variation, and disease. *Arch Biochem Biophys* 389:84–93, 2001
64. FUKAI T, FOLZ R, LANDMESSER U, HARRISON D: Extracellular superoxide dismutase and cardiovascular disease. *Cardiovasc Res* 55:239–249, 2002
65. BALIGA R, UEDA N, WALKER P, SHAH S: Oxidant mechanisms in toxic acute renal failure. *Drug Metab Rev* 31:971–997, 1999
66. MCDONALD J, GALLEY H, WEBSTER N: Oxidative stress and gene expression in sepsis. *Br J Anaesth* 90:221–232, 2003
67. DRUML W, SCHWARZENHOFER M, APSNER R, HORL W: Fat-soluble vitamins in patients with acute renal failure. *Miner Electrolyte Metab* 24:220–226, 1998
68. METNITZ P, FISCHER M, BARTENS C, et al: Impact of acute renal failure on antioxidant status in multiple organ failure. *Acta Anaesthesiol Scand* 44:236–240, 2000
69. BALAKRISHNAN V, BLUMBERG J, PEREIRA BJ, JABER BL: Antioxidant and oxidative stress indices in dialysis-dependent acute renal failure. *Blood Purif* 21:213–219, 2003
70. STORY A, RONCO C, BELLOMO R: Trace element and vitamin concentrations and losses in critically ill patients treated with continuous venovenous hemofiltration. *Crit Care Med* 27:220–223, 1999
71. YSEBAERT D, DE GREEF K, VERCAUTEREN S, et al: Identification and kinetics of leukocytes after severe ischaemia/reperfusion renal injury. *Nephrol Dial Transplant* 15:1562–1574, 2000
72. ALONSO A, LAU J, JABER B, et al: Prevention of radiocontrast nephropathy with N-acetylcysteine in patients with chronic kidney disease: A meta-analysis of randomized, controlled trials. *Am J Kidney Dis* 43:1–9, 2004
73. CONESA E, VALERO F, NADAL J, et al: N-acetyl-L-cysteine improves renal medullary hypoperfusion in acute renal failure. *Am J Physiol Regul Integr Comp Physiol* 281:R730–R737, 2001
74. HEYMAN S, GOLDFARB M, SHINA A, et al: N-acetylcysteine ameliorates renal microcirculation: studies in rats. *Kidney Int* 63:634–641, 2003
75. WADDINGTON S, MOSLEY K, CATTELL V: Induced nitric oxide (NO) synthesis in heterologous nephrotoxic nephritis; effects of selective inhibition in neutrophil-dependent glomerulonephritis. *Clin Exp Immunol* 118:309–314, 1999
76. PARK K, BYUN J, KRAMERS C, et al: Inducible nitric-oxide synthase is an important contributor to prolonged protective effects of ischemic preconditioning in the mouse kidney. *J Biol Chem* 278:27256–27266, 2003
77. YAMASHITA J, OGATA M, ITOH M, et al: Role of nitric oxide in the renal protective effects of ischemic preconditioning. *J Cardiovasc Pharmacol* 42:419–427, 2003
78. SCHRAMM L, LA M, HEIDBREDE E, et al: L-arginine deficiency and supplementation in experimental acute renal failure and in human kidney transplantation. *Kidney Int* 61:1423–1432, 2002
79. WILSON AG, DE VRIES N, POCIOT F, et al: An allelic polymorphism within the human tumor necrosis factor- α promoter region is strongly associated with HLA A1, B8, and DR3 alleles. *J Exp Med* 177:557–560, 1993
80. MORSE H, OLOMOLAIYE O, WOOD N, et al: Induced heteroduplex genotyping of TNF- α , IL-1 β , IL-6 and IL-10 polymorphisms associated with transcriptional regulation. *Cytokine* 11:789–795, 1999
81. ABRAHAM LJ, KROEGER KM: Impact of the -308 TNF promoter polymorphism on the transcriptional regulation of the TNF gene: relevance to disease. *J Leukoc Biol* 66:562–566, 1999
82. WILSON AG, SYMONS JA, McDOWELL TL, et al: Effects of a polymorphism in the human tumor necrosis factor [alpha] promoter on transcriptional activation. *Proc Natl Acad Sci* 94:3195–3199, 1997
83. KROEGER KM, CARVILLE KS, ABRAHAM LJ: The -308 tumor necrosis factor-[alpha] promoter polymorphism effects transcription. *Mol Immunol* 34:391–399, 1997
84. WARZUCHA K, RIBEIRO P, BIENVENU J: Genetic polymorphisms in the tumor necrosis factor locus influence non-Hodgkin's lymphoma outcome. *Blood* 91:3574–3581, 1998
85. LOUIS E, FRANCHIMONT D, PIRON A: Tumor necrosis factor gene polymorphism influences TNF- α production in lipopolysaccharide-stimulated whole blood cell culture in healthy humans. *Clin Exp Immunol* 113:401–406, 1998
86. ABRAHAM L, DU D, ZAHEDI K, et al: Haplotypic polymorphisms of the TNFB gene. *Immunogenetics* 33:50–53, 1991
87. POCIOT F, MOLVIG J, WOGENSEN L, et al: A tumour necrosis factor beta gene polymorphism in relation to monokine secretion and insulin-dependent diabetes mellitus. *Scand J Immunol* 33:37–49, 1991
88. TABRIZI A, ZEHNBAUER B, FREEMAN B, BUCHMAN T: Genetic markers in sepsis. *J Am Coll Surg* 192:106–117, 2001
89. HEESSEN M, KUNZ D, BACHMANN-MENNENGA B, et al: Linkage disequilibrium between tumor necrosis factor (TNF)- α -308 G/A promoter and TNF- β NcoI polymorphisms: Association with TNF- α response of granulocytes to endotoxin stimulation. *Crit Care Med* 31:211–214, 2003
90. MIRA J-P, CARIU A, GRALL F, et al: Association of TNF2, a TNF-[alpha] promoter polymorphism, with septic shock susceptibility and mortality: A multicenter study. *JAMA* 282:561–568, 1999
91. TANG GJ, HUANG SL, YIEN HW, et al: Tumor necrosis factor gene polymorphism and septic shock in surgical infection. *Crit Care Med* 28:2733–2736, 2000
92. WEITKAMP JH, STUBER F, BARTMANN P: Pilot study assessing TNF

- gene polymorphism as a prognostic marker for disease progression in neonates with sepsis. *Infection* 28:92–96, 2000
93. WUNDERINK RG, WATERER GW, CANTOR RM, QUASNEY MW: Tumor necrosis factor gene polymorphisms and the variable presentation and outcome of community-acquired pneumonia. *Chest* 121:87S, 2002
 94. NADEL S, NEWPORT MJ, BOOY R, LEVIN M: Variation in the tumor necrosis factor- α promoter region may be associated with death from meningococcal disease. *J Infect Dis* 174:878–880, 1996
 95. MCGUIRE W, HILL AV, ALLSOPP CE, et al: Variation of the TNF- α promoter region associated with susceptibility to cerebral malaria. *Nature* 371:508–510, 1994
 96. CABRERA M, SHAW MA, SHARPLES C: Polymorphism in tumor necrosis factor genes associated with mucocutaneous leishmaniasis. *J Exp Med* 182:1259–1264, 1995
 97. STUBER F, PETERSEN M, BOKELMANN F, SCHADE U: A genomic polymorphism within the tumor necrosis factor locus influences plasma tumor necrosis factor- α concentrations and outcome of patients with severe sepsis. *Crit Care Med* 24:381–384, 1996
 98. MAJETSCHAK M, FLOHE S, OBERTACKE U, et al: Relation of a TNF gene polymorphism to severe sepsis in trauma patients. *Ann Surg* 230:207–214, 1999
 99. YENDE S, QUASNEY MW, TOLLEY E, et al: Association of tumor necrosis factor gene polymorphisms and prolonged mechanical ventilation after coronary artery bypass surgery. *Crit Care Med* 31:133–140, 2003
 100. SCHROEDER S, BORGER N, WRIGGE H, et al: A tumor necrosis factor gene polymorphism influences the inflammatory response after cardiac operation. *Ann Thorac Surg* 75:534–537, 2003
 101. YENDE S, QUASNEY M, ZHANG Q, et al: Impact of cytokine gene polymorphisms on outcomes of coronary artery bypass graft surgery. *Chest* 121:86S, 2002
 102. SHAIKH K, YENDE S, WATERER GW, et al: Effect of tumor necrosis factor gene polymorphisms on restoration of body temperature after coronary artery bypass graft surgery. *Chest* 118:109S, 2000
 103. JABER B, RAO M, GUO D, et al: Cytokine promoter gene polymorphisms and mortality in acute renal failure. *Cytokine* 25:212–219, 2004
 104. DOMINICI R, CATTANEO M, Malferrari G, et al: Cloning and functional analysis of the allelic polymorphism in the transcription regulatory region of interleukin-1 α . *Immunogenetics* 54:82–86, 2002
 105. READ R, CANNINGS C, NAYLOR S, et al: Variation within genes encoding interleukin-1 and the interleukin-1 receptor antagonist influence the severity of meningococcal disease. *Ann Intern Med* 138:534–541, 2003
 106. VIRTA M, HURME M, HELMINEN M: Increased frequency of interleukin-1 β (–511) allele 2 in febrile seizures. *Pediatr Neurol* 26:192–195, 2002
 107. BERGER P, MCCONNELL J, NUNN M, et al: C-reactive protein levels are influenced by common IL-1 gene variations. *Cytokine* 17:171–174, 2002
 108. COX A, CAMP N, NICKLIN M, et al: An analysis of linkage disequilibrium in the interleukin-1 gene cluster, using a novel grouping method for multiallelic markers. *Am J Hum Genet* 62:1180–1188, 1998
 109. SANTTILA S, SAVINAINEN K, HURME M: Presence of the IL-1RA allele 2 (IL1RN*2) is associated with enhanced IL-1 β production in vitro. *Scand J Immunol* 47:195–198, 1998
 110. ANDUS T, DAIG R, VOGL D, et al: Imbalance of the interleukin 1 system in colonic mucosa—Association with intestinal inflammation and interleukin 1 receptor antagonist [corrected] genotype 2. *Gut* 41:651–657, 1997
 111. TOUNTAS N, CASINI-RAGGI V, YANG H, et al: Functional and ethnic association of allele 2 of the interleukin-1 receptor antagonist gene in ulcerative colitis. *Gastroenterology* 117:806–813, 1999
 112. VAMVAKOPOULOS J, GREEN C, METCALFE S: Genetic control of IL-1 β bioactivity through differential regulation of the IL-1 receptor antagonist. *Eur J Immunol* 32:2988–2996, 2002
 113. FANG X, SCHRODER S, HOEF A, STUBER F: Comparison of two polymorphisms of the interleukin-1 gene family: Interleukin-1 receptor antagonist polymorphism contributes to susceptibility to severe sepsis. *Crit Care Med* 27:1330–1334, 1999
 114. MA P, CHEN D, PAN J, DU B: Genomic polymorphism within interleukin-1 family cytokines influences the outcome of septic patients. *Crit Care Med* 30:1046–1050, 2002
 115. WILKINSON R, PATEL P, LLEWELYN M, et al: Influence of polymorphism in the genes for the interleukin (IL)-1 receptor antagonist and IL-1 β on tuberculosis. *J Exp Med* 189:1863–1874, 1999
 116. AMOLI M, THOMSON W, HAJEER A, et al: Interleukin 1 receptor antagonist gene polymorphism is associated with severe renal involvement and renal sequelae in Henoch-Schönlein purpura. *J Rheumatol* 29:1404–1407, 2002
 117. RAY K, CAMP N, BENNETT C, et al: Genetic variation at the interleukin-1 locus is a determinant of changes in soluble endothelial factors in patients with acute coronary syndromes. *Clin Sci (Lond)* 103:303–310, 2002
 118. BORGMANN S, ENDISCH G, HACKER U, et al: Proinflammatory genotype of interleukin-1 and interleukin-1 receptor antagonist is associated with ESRD in proteinase 3–ANCA vasculitis patients. *Am J Kidney Dis* 41:933–942, 2003
 119. BARTON K: IL-6: insights into novel biological activities. *Clin Immunol Immunopathol* 85:16–20, 1997
 120. FISHMAN D, FAULDS G, JEFFERY R, et al: The effect of novel polymorphisms in the interleukin-6 (IL-6) gene on IL-6 transcription and plasma IL-6 levels, and an association with systemic-onset juvenile chronic arthritis. *J Clin Invest* 102:1369–1376, 1998
 121. FERRARI S, AHN-LUONG L, GARNERO P, et al: Two promoter polymorphisms regulating interleukin-6 gene expression are associated with circulating levels of C-reactive protein and markers of bone resorption in postmenopausal women. *J Clin Endocrinol Metab* 88:255–259, 2003
 122. FERNANDEZ-REAL JM, BROCH M, VENDRELL J, et al: Interleukin-6 gene polymorphism and lipid abnormalities in healthy subjects. *J Clin Endocrinol Metab* 85:1331–1333, 2000
 123. COX ED HS, DiMERCURIO BS, WESLEY RA, et al: Cytokine polymorphic analyses indicate ethnic differences in the allelic distribution of interleukin-2 and interleukin-6. *Transplantation* 72:720–726, 2001
 124. HEESEN M, OBERTACKE U, SCHADE F, et al: The interleukin-6 G(–174)C polymorphism and the ex vivo interleukin-6 response to endotoxin in severely injured blunt trauma patients. *Eur Cytokine New* 13:72–77, 2002
 125. SCHLUTER B, RAUFHAK C, ERREN M, et al: Effect of the interleukin-6 promoter polymorphism (–174 G/C) on the incidence and outcome of sepsis. *Crit Care Med* 30:32–37, 2002
 126. GALLAGHER P, LOWE G, FITZGERALD T, et al: Association of IL-10 polymorphism with severity of illness in community acquired pneumonia. *Thorax* 58:154–156, 2003
 127. HAHN A, KASTEN-JOLLY J, CONSTANTINO D, et al: TNF- α , IL-6, IFN- γ , and IL-10 gene expression polymorphisms and the IL-4 receptor α -chain variant Q576R: effects on renal allograft outcome. *Transplantation* 72:660–665, 2001
 128. BRULL DJ, MONTGOMERY HE, SANDERS J, et al: Interleukin-6 gene –174g>c and –572g>c promoter polymorphisms are strong predictors of plasma interleukin-6 levels after coronary artery bypass surgery. *Arterioscler Thromb Vasc Biol* 21:1458–1463, 2001
 129. BURZOTTA F, IACOVIELLO L, Di CASTELNUOVO A, et al: Relation of the –174 G/C polymorphism of interleukin-6 to interleukin-6 plasma levels and to length of hospitalization after surgical coronary revascularization. *Am J Cardiol* 88:1125–1128, 2001
 130. TRESZL A, TOTHEYN P, KOCIS I, et al: Interleukin genetic variants and the risk of renal failure in infants with infection. *Pediatr Nephrol* 17:713–717, 2002
 131. HOMS E, RIBEIRO-ALVES M, LOPES DE FARIA J, DIAS E: Interleukin-6 stimulates tubular regeneration in rats with glycerol-induced acute renal failure. *Nephron* 92:192–199, 2002
 132. WESTENDORP RG LJ, HUIZINGA TW, ELOUALI AH, et al: Genetic influence on cytokine production and fatal meningococcal disease. *Lancet* 349:170–173, 1997
 133. CRAWLEY E, KAY R, SILLIBOURNE J, et al: Polymorphic haplotypes of the interleukin-10 5' flanking region determine variable interleukin-10 transcription and are associated with particular phenotypes of juvenile rheumatoid arthritis. *Arthritis Rheum* 42:1101–1108, 1999

134. VAN DER POL WLHT, VIDARSSON G, VAN DER LINDEN MW, *et al*: Relevance of Fcγ receptor and interleukin-10 polymorphisms for meningococcal disease. *J Infect Dis* 184:1548–1555, 2001
135. PELLETIER R, PRAVICA V, PERREY C, *et al*: Evidence for a genetic predisposition towards acute rejection after kidney and simultaneous kidney-pancreas transplantation. *Transplantation* 70:674–680, 2000
136. GIRNDT M, SESTER U, SESTER M, *et al*: Impaired cellular immune function in patients with end-stage renal failure. *Nephrol Dial Transplant* 14:2807–2810, 1999
137. LOWE PR, GALLEY HF, ABDEL-FATTAH A, WEBSTER NR: Influence of interleukin-10 polymorphisms on interleukin-10 expression and survival in critically ill patients. *Crit Care Med* 31:34–38, 2003
138. HOLDSWORTH S, KITCHING A, TIPPING P: Chemokines as therapeutic targets in renal disease. *Curr Opin Nephrol Hypertens* 9:505–511, 2000
139. MODI W, DEAN M, SEUANEZ H, *et al*: Monocyte-derived neutrophil chemotactic factor (MDNCF/IL-8) resides in a gene cluster along with several other members of the platelet factor 4 gene superfamily. *Hum Genet* 84:185–187, 1990
140. JIANG Z, OKHUYSEN P, GUO D, *et al*: Genetic susceptibility to enteroaggregative *Escherichia coli* diarrhea: Polymorphism in the IL-8 promoter region at position –251. *J Infect Dis* 188:506–511, 2003
141. HULL J, THOMSON A, KWIATKOWSKI D: Association of respiratory syncytial virus bronchiolitis with the interleukin 8 gene region in UK families. *Thorax* 55:1023–1027, 2000
142. YU Y, CHADEE K: The 3'-untranslated region of human interleukin-8 mRNA suppresses IL-8 gene expression. *Immunology* 102:498–505, 2001
143. AMOLI MM, THOMSON W, HAJEER AH, *et al*: Interleukin 8 gene polymorphism is associated with increased risk of nephritis in cutaneous vasculitis. *J Rheumatol* 29:2367–2370, 2002
144. ROVIN B, LU L, ZHANG X: A novel interleukin-8 polymorphism is associated with severe systemic lupus erythematosus nephritis. *Kidney Int* 62:261–265, 2002
145. ROVIN BH, LU L, SAXENA R: A novel polymorphism in the MCP-1 gene regulatory region that influences MCP-1 expression. *Biochem Biophys Res Commun* 259:344–348, 1999
146. KIM H, LEE D, YANG S, *et al*: The polymorphism of monocyte chemoattractant protein-1 is associated with the renal disease of SLE. *Am J Kidney Dis* 40:1146–1152, 2002
147. KRUGER B, SCHROPP B, ASHKAN R, *et al*: A Monocyte chemoattractant protein-1 (MCP-1) polymorphism and outcome after renal transplantation. *J Am Soc Nephrol* 13:2585–2589, 2002
148. POLTORAK A, SMIRNOVA I, HE X, *et al*: Genetic and physical mapping of the Lps locus: Identification of the toll-4 receptor as a candidate gene in the critical region. *Blood Cells Mol Dis* 24:340–355, 1998
149. POLTORAK A, HE X, SMIRNOVA I, *et al*: Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: Mutations in Tlr4 gene. *Science* 282:2085–2088, 1998
150. HOSHINO K, TAKEUCHI O, KAWAI T, *et al*: Cutting edge: Toll-like receptor 4 (TLR4)-deficient mice are hyporesponsive to lipopolysaccharide: Evidence for TLR4 as the Lps gene product. *J Immunol* 162:3749–3752, 1999
151. BIHL F, SALEZ L, BEAUBIER M, *et al*: Overexpression of toll-like receptor 4 amplifies the host response to lipopolysaccharide and provides a survival advantage in transgenic mice. *J Immunol* 170:6141–6150, 2003
152. ARBOUR N, LORENZ E, SCHUTTE B, *et al*: TLR4 mutations are associated with endotoxin hyporesponsiveness in humans. *Nat Genet* 25:187–191, 2000
153. CHILD N, YANG I, PULLETT M, *et al*: Polymorphisms in toll-like receptor 4 and the systemic inflammatory response syndrome. *Biochem Soc Trans* 31:652–653, 2003
154. AGNESE D, CALVANO J, HAHM S, *et al*: Human toll-like receptor 4 mutations but not CD14 polymorphisms are associated with an increased risk of gram-negative infections. *J Infect Dis* 186:1522–1525, 2002
155. LORENZ E, MIRA J, CORNISH K, *et al*: A novel polymorphism in the toll-like receptor 2 gene and its potential association with staphylococcal infection. *Infect Immunol* 68:6398–6401, 2000
156. LORENZ E, MIRA J, FREES K, SCHWARTZ D: Relevance of mutations in the TLR4 receptor in patients with gram-negative septic shock. *Arch Intern Med* 162:1028–1032, 2002
157. BOCHUD P, HAWN T, ADEREM A: Cutting edge: A toll-like receptor 2 polymorphism that is associated with lepromatous leprosy is unable to mediate mycobacterial signaling. *J Immunol* 170:3451–3454, 2003
158. VILLAR J, MENDEZ-ALVAREZ S: Heat shock proteins and ventilator-induced lung injury. *Curr Opin Crit Care* 9:9–14, 2003
159. FEKETE A, TRESZL A, TOTTH-HEYEN P, *et al*: Association between heat shock protein 72 gene polymorphism and acute renal failure in premature neonates. *Pediatr Res* 54:452–455, 2003
160. SCHRODER O, SCHULTE K-M, OSTERMANN P, *et al*: Heat shock protein 70 genotypes HSPA1B and HSPA1L influence cytokine concentrations and interfere with outcome after major injury. *Crit Care Med* 31:76–79, 2003
161. ASEA A, KRAEFT S, KURT-JONES E, *et al*: HSP70 stimulates cytokine production through a CD14-dependant pathway, demonstrating its dual role as a chaperone and cytokine. *Nat Med* 6:435–442, 2000
162. KELLY K: Stress response proteins and renal ischemia. *Minerva Urol Nefrol* 54:81–91, 2002
163. WATERER G, ELBAHLAWAN L, QUASNEY M, *et al*: Heat shock protein 70–2+1267 AA homozygotes have an increased risk of septic shock in adults with community-acquired pneumonia. *Crit Care Med* 31:1367–1372, 2003
164. NATH K: Tubulointerstitial changes as a major determinant in the progression of renal damage. *Am J Kid Dis* 20:1–17, 1992
165. SHIRAIISHI F, CURTIS L, TRUONG L, *et al*: Heme oxygenase-1 gene ablation or expression modulates cisplatin-induced renal tubular apoptosis. *Am J Physiol Renal Physiol* 278:F726–F736, 2000
166. BLYDT-HANSEN T, KATORI M, LASSMAN C, *et al*: Gene transfer-induced local heme oxygenase-1 overexpression protects rat kidney transplants from ischemia/reperfusion injury. *J Am Soc Nephrol* 14:745–754, 2003
167. POSS K, TONEGAWA S: Reduced stress defense in heme oxygenase deficient cells. *Proc Natl Acad Sci USA* 94:10925–10930, 1997
168. YACHIE A, NIIDA Y, WADA T, *et al*: Oxidative stress causes enhanced endothelial cell injury in human heme oxygenase-1 deficiency. *J Clin Invest* 103:129–135, 1999
169. OTTERBEIN L, BACH F, ALAM J, *et al*: Carbon monoxide has anti-inflammatory effects involving the mitogen-activated protein kinase pathway. *Nat Med* 6:422–428, 2000
170. LEE T-S, CHAU L-Y: Heme oxygenase-1 mediates the anti-inflammatory effect of interleukin-10 in mice. *Nat Med* 8:240–246, 2002
171. KUTTY R, KUTTY G, RODRIGUES I, *et al*: Chromosomal localization of the human heme oxygenase genes: heme oxygenase-1 (HMOX 1) maps to chromosome 22q12 and heme oxygenase-2 (HMOX 2) maps to chromosome 16p13.3. *Genomics* 20:513–516, 1994
172. LAVROVSKY Y, SCHWARTZMAN M, LEVERE R, *et al*: Identification of binding sites for transcription factors NF-κB and AP-2 in the promoter region of the human heme oxygenase-1 gene. *Proc Natl Acad Sci USA* 91:5987–5991, 1994
173. CHEN Y-H, LIN S, LIN M, *et al*: Microsatellite polymorphism in promoter of heme oxygenase-1 gene is associated with susceptibility to coronary artery disease in type 2 diabetic patients. *Hum Genet* 111:1–8, 2002
174. EXNER M, SCHILLINGER M, MINAR E, *et al*: Heme oxygenase-1 gene promoter microsatellite polymorphism is associated with restenosis after percutaneous transluminal angioplasty. *J Endovasc Ther* 8:433–440, 2001
175. SCHILLINGER M, EXNER M, MLEKUSCH W, *et al*: Heme oxygenase-1 gene promoter polymorphism is associated with abdominal aortic aneurysm. *Thromb Res* 106:131–136, 2002
176. YAMADA N, YAMAYA M, OKINAGA S, *et al*: Microsatellite polymorphism in the heme oxygenase-1 gene promoter is associated with susceptibility to emphysema. *Am J Hum Genet* 66:187–195, 2000
177. HAYES J, STRANGE R: Potential contribution of the glutathione S-transferase supergene family to resistance to oxidative stress. *Free Radic Res* 22:193, 1995
178. INOUE N, KAWASHIMA S, KANAZAWA K, *et al*: Polymorphism of the NADH/NADPH oxidase p22phox gene in patients with coronary artery disease. *Circulation* 97:135–137, 1998

179. GUZIK T, WEST N, BLACK E, *et al*: Functional effect of the C242T polymorphism in the NAD(P)H oxidase p22phox gene on vascular superoxide production in atherosclerosis. *Circulation* 102:1744–1747, 2000
180. ZAFARI A, DAVIDOFF M, AUSTIN H, *et al*: The A640G and C242T p22(phox) polymorphisms in patients with coronary artery disease. *Antioxid Redox Signal* 4:675–680, 2002
181. SCHNEIDER M, HILGERS K, HUANG Y, *et al*: The C242T p22phox polymorphism and endothelium-dependent vasodilation in subjects with hypercholesterolaemia. *Clin Sci (Lond)* 105:97–103, 2003
182. STANGER O, RENNER W, KHOSCHSORUR G, *et al*: NADH/NADPH oxidase p22 phox C242T polymorphism and lipid peroxidation in coronary artery disease. *Clin Physiol* 21:718–722, 2001
183. NAKANO T, MATSUNAGA S, NAGATA A, MARUYAMA T: NAD(P)H oxidase p22phox Gene C242T polymorphism and lipoprotein oxidation. *Clin Chim Acta* 335:101–107, 2003
184. WOLF G, PANZER U, HARENDZA S, *et al*: No association between a genetic variant of the p22(phox) component of NAD(P)H oxidase and the incidence and progression of IgA nephropathy. *Nephrol Dial Transplant* 17:1509–1512, 2002
185. RAJMAKERS M, ROES E, STEEGERS E, PETERS W: The C242T-polymorphism of the NADPH/NADH oxidase gene p22phox subunit is not associated with pre-eclampsia. *J Hum Hypertens* 16:423–425, 2002
186. PIEDRAFITA F, MOLANDER R, VANSANT G, *et al*: An Alu element in the myeloperoxidase promoter contains a composite SP1-thyroid hormone-retinoic acid response element. *J Biol Chem* 271:14412–14420, 1996
187. PECOITS-FILHO R, STENVINKEL P, MARCHLEWSKA A, *et al*: A functional variant of the myeloperoxidase gene is associated with cardiovascular disease in end-stage renal disease patients. *Kidney Int (Suppl 84)*:S172–S176, 2003
188. SANDSTROM J, NILSSON P, KARLSSON K, MARKLUND S: 10-fold increase in human plasma extracellular superoxide dismutase content caused by a mutation in heparin-binding domain. *J Biol Chem* 269:19163–19166, 1994
189. FORSBERG L, LYRENÄS L, DE FAIRE U, MORGENSTERN R: A common functional C-T substitution polymorphism in the promoter region of the human catalase gene influences transcription factor binding, reporter gene transcription and is correlated to blood catalase levels. *Free Radic Biol Med* 30:500–505, 2001
190. SHEN Q, TOWNES P, PADDEN C, NEWBURGER P: An in-frame trinucleotide repeat in the coding region of the human cellular glutathione peroxidase (GPX1) gene: In vivo polymorphism and in vitro instability. *Genomics* 23:292–294, 1994
191. MOSCOW J, MORROW C, HE R, *et al*: Structure and function of the 5'-flanking sequence of the human cytosolic selenium-dependent glutathione peroxidase gene (hgp1). *J Biol Chem* 267:5949–5958, 1994
192. FORSBERG L, DE FAIRE U, MARKLUND S, *et al*: Phenotype determination of a common Pro-Leu polymorphism in human glutathione peroxidase 1. *Blood Cells Mol Dis* 26:423–426, 2000
193. SEIDEGARD J, VORACHEK W, PERO R, PEARSON W: Hereditary differences in the expression of the human glutathione transferase active on trans-stilbene oxide are due to a gene deletion. *Proc Natl Acad Sci USA* 85:7293–7297, 1988
194. BOARD P: Biochemical genetics of glutathione-S-transferase in man. *Am J Hum Genet* 33:36–43, 1981
195. WIDERSTEN M, PEARSON W, ENGSTROM A, MANNERVIK B: Heterologous expression of the allelic variant mu-class glutathione transferases mu and psi. *Biochem J* 276:519–524, 1991
196. SEIDEGARD J, PERO R, MARKOWITZ M, *et al*: Isoenzyme(s) of glutathione transferase (class Mu) as a marker for the susceptibility to lung cancer: A follow up study. *Carcinogenesis* 11:33–36, 1990
197. ST PETER S, IMBER C, JONES D, *et al*: Genetic determinants of delayed graft function after kidney transplantation. *Transplantation* 74:809–813, 2002
198. HOBBS M, UDHAYAKUMAR V, LEVESQUE M, *et al*: A new NOS2 promoter polymorphism associated with increased nitric oxide production and protection from severe malaria in Tanzanian and Kenyan children. *Lancet* 360:1468–1475, 2002
199. NABEL E: Genomic medicine: Cardiovascular disease. *N Engl J Med* 349:60–72, 2003
200. RIDKER P, STAMPFER M: Assessment of genetic markers for coronary thrombosis: Promise and precaution. *Lancet* 353:687–688, 1999
201. CONSORTIUM TIH: The International HapMap Project. *Nature* 426:789–796, 2003
202. NEWMAN TB, BROWNER WS, HULLEY SB: Enhancing causal inference in observational studies, in *Designing Clinical Research*, edited by Hulley SB, Cummings SR, Baltimore, Williams & Wilkins, 1988, pp 98–109
203. ROSENTHAL N, SCHWARTZ R: In search of perverse polymorphisms. *N Engl J Med* 338:122–124, 1998
204. STUBER F: Another definite candidate gene for genetic predisposition of sepsis: Interleukin-10. *Crit Care Med* 31:314–315, 2003
205. JABER BL, LIANGOS O, PEREIRA B, BALAKRISHNAN V: Polymorphism of immuno-modulatory cytokine genes: Implications in acute renal failure. *Blood Purif* 22:101–111, 2004
206. BAILLY S, DI GIOVINE F, DUFF G: Polymorphic tandem repeat region in interleukin-1 alpha intron 6. *Hum Genet* 91:85–86, 1993
207. POCIOT F, MOLVIG J, WOGENSEN L, *et al*: A TaqI polymorphism in the human interleukin-1 beta (IL-1 beta) gene correlates with IL-1 beta secretion in vitro. *Eur J Clin Invest* 22:396–402, 1992
208. KANG T, LEE S, CHAE G: A polymorphism in the toll-like receptor 2 is associated with IL-12 production from monocyte in lepromatous leprosy. *Cytokine* 20:56–62, 2002
209. MCGUIRE W, KNIGHT J, HILL A, *et al*: Severe malarial anemia and cerebral malaria are associated with different tumor necrosis factor promoter alleles. *J Infect Dis* 179:287–290, 1999
210. ZHANG D, LI J, JIANG Z, *et al*: Association of two polymorphisms of tumor necrosis factor gene with acute severe pancreatitis. *J Surg Res* 112:138–143, 2003
211. ASENSI V, ALVAREZ V, VALLE E, *et al*: IL-1alpha (–889) promoter polymorphism is a risk factor for osteomyelitis. *Am J Med Genet* 119A:132–136, 2003
212. READ R, CAMP N, DI GIOVINE F, *et al*: An interleukin-1 genotype is associated with fatal outcome of meningococcal disease. *J Infect Dis* 182:1557–1560, 2000
213. DELGADO J, BAENA A, THIM S, GOLDFELD A: Ethnic-specific genetic associations with pulmonary tuberculosis. *J Infect Dis* 186:1463–1468, 2002
214. LIN M, STORER B, MARTIN P, *et al*: Relation of an interleukin-10 promoter polymorphism to graft-versus-host disease and survival after hematopoietic-cell transplantation. *N Engl J Med* 349:2201–2210, 2003